

THE FROBENIUS THEOREM ON N -DIMENSIONAL QUANTUM HYPERPLANE

C. CIUPALĂ

ABSTRACT. In this paper we introduce the universal ρ -differential calculus on a ρ -algebra and we prove the universality of the construction. We also present submanifolds, distributions, linear connections and two different ρ -differential calculi on S_N^q : the algebra of forms and the algebra of universal differential forms of S_N^q . Finally we prove the Frobenius theorem on the N -dimensional quantum hyperplane S_N^q in a general case, for any ρ -differential calculus on S_N^q .

1. INTRODUCTION

The basic idea of noncommutative geometry is to replace an algebra of smooth functions defined on a smooth manifold by an abstract associative algebra A which is not necessarily commutative. In the context of noncommutative geometry the basic role is the generalization of the notion of differential forms (on a manifold). A (noncommutative) differential calculus over the associative algebra A is a \mathbb{Z} -graded algebra $\Omega(A) = \bigoplus_{n \geq 0} \Omega^n(A)$ (where $\Omega^n(A)$ are A -bimodules and $\Omega^0(A) = A$) together with a linear operator $d : \Omega^n(A) \rightarrow \Omega^{n+1}(A)$ satisfying $d^2 = 0$ and $d(\omega\omega') = (d\omega)\omega' + (-1)^n\omega d\omega'$ where $\omega \in \Omega^n(A)$.

There are studied some differential calculi associated to A , here we recall two of them: the algebra of forms of A in [8] and the algebra of universal differential forms of A in [2]. These two differential calculi are studied even in the case when A is a superalgebra in [10] and in [11], but when A is a ρ -algebra there is studied only the algebra of forms of A in [1].

Received March 3, 2004.

2000 *Mathematics Subject Classification.* Primary 81R60, 16W99, 53C04.

Key words and phrases. noncommutative geometry, linear connections, submanifolds, distributions, quantum hyperplane.

In this paper we introduce the notion of universal derivation of the order (α, β) , we construct the algebra of universal differential forms of a ρ -algebra A and we prove the universality of the construction. We apply this to the particular case of the N -dimensional quantum hyperplane S_N^q and thus we obtain a new differential calculus $\Omega_\alpha(S_N^q)$ on S_N^q , denoted the algebra of universal differential forms of S_N^q .

On the other hand we present submanifolds algebra on S_N^q in the context of noncommutative geometry, we also define linear connections, distributions on S_N^q and we prove the Frobenius theorem on S_N^q , for any ρ -differential calculus ΩS_N^q on S_N^q .

2. DIFFERENTIAL CALCULUS ON ρ -ALGEBRAS

In this section we define the differential calculus on a ρ -algebra A and we give two examples of such differential calculus on A : the algebra of forms $\Omega(A)$ of a ρ -commutative algebra A from [1] and the algebra of universal differential forms $\Omega_\alpha(A)$ of a ρ -algebra A . We briefly review the basic notions of ρ -algebras and ρ -modules on ρ -algebras. (see [1], [3], [5] and [6] for details).

First we review the basic notions about ρ -algebras, ρ -bimodules over a ρ -algebra and the derivations on ρ -bimodules.

Let G be an abelian group, additively written, and let A be a G -graded algebra. A is a ρ -algebra if there is a map $\rho : G \times G \rightarrow k$ which satisfies the relations

$$(1) \quad \rho(a, b) = \rho(b, a)^{-1} \quad \text{and} \quad \rho(a + b, c) = \rho(a, c)\rho(b, c), \quad \text{for any } a, b, c \in G.$$

The G -degree of a (nonzero) homogeneous element f of A is denoted by $|f|$. The ρ -algebra A is ρ -commutative if $fg = \rho(|f|, |g|)gf$ for any $f \in A_{|f|}$ and $g \in A_{|g|}$.

The morphism $f : M \rightarrow N$ between the ρ -bimodules M and N over the ρ -algebra A is of degree $\beta \in G$ if $f : M_\alpha \rightarrow N_{\alpha+\beta}$ for any $\alpha \in \dot{G}$, $f(am) = \rho(\alpha, |a|)af(m)$ and $f(ma) = f(m)a$ for any $a \in A_{|a|}$ and $m \in M$.

Definition 1. Let $\alpha, \beta \in G$ and M a ρ -bimodule over the ρ -algebra A . A ρ -derivation of order (α, β) on M (ρ -derivation of degree α and of G -degree β) is a linear map $X : A \rightarrow M$, which fulfils the properties:

1. $X : A_* \rightarrow M_{*+\beta}$,
2. $X(fg) = (Xf)g + \rho(\alpha, |f|)f(Xg)$, for any $f \in A_{|f|}$ and $g \in A$.

The left product $fX : A \rightarrow M$ between the element $f \in A$ and a ρ -derivation X of the order (α, β) is defined in a natural way: by $(fX)(g) = fX(g)$, for any $g \in A$.

Proposition 1. *If is X a ρ -derivation of order (α, β) on M and $f \in A_{|f|}$, then fX is a derivation of the order $(|f| + \alpha, |f| + \beta)$ on M if and only if the algebra A is ρ -commutative.*

Proof. We have to show that $(fX)(gh) = ((fX)g)h + \rho(|f| + \alpha, |g|)g(fX)h$ and $fX : A_* \rightarrow M_{*+|f|+|X|}$, for any $h \in A_{|h|}$ and $f \in A_{|f|}$.

$$\begin{aligned} (fX)(gh) &= f(X(gh)) = f(X(g)h + \rho(\alpha, |g|)gX(h)) \\ &= (fX)(g)h + \rho(\alpha, |g|)fgX(h) = (fX)(g)h + \rho(\alpha, |g|)\rho(|f|, |g|)gfX(h) \\ &= (fX)(g)h + \rho(\alpha, |f| + |g|)g(fX)(h). \end{aligned}$$

The second relation is obvious. □

The linear map $X : A \rightarrow A$ is a ρ -derivation on A if it is a ρ -derivation of the order $(|X|, |X|)$. Next we denote by $\rho\text{-Der}(A)$ the ρ -bimodule over A of the all ρ -derivations on A .

Definition 2. Let A be a ρ -algebra. We say that $\Omega(A) = \bigoplus_{(n,\beta) \in \mathbb{Z} \times G} \Omega_\beta^n(A)$ is a ρ -differential calculus on A if $\Omega(A)$ is a $\mathbb{Z} \times G$ -graded algebra, $\Omega(A)$ is a ρ -bimodule over A , $\Omega^0(A) = \bigoplus_{\beta \in G} \Omega_\beta^0(A) = A$ and there is an element $\alpha \in G$ and ρ -derivation $d : \Omega(A) \rightarrow \Omega(A)$ of the order $((1, \alpha), (1, 0))$ such that $d^2 = 0$.

The first example of a ρ -differential calculus over the ρ -commutative algebra A is the algebra of forms $(\Omega(A), d)$ of A from [1].

The second example of a ρ -differential calculus over a ρ -algebra is the universal differential calculus of A from the next subsection.

2.1. The algebra of universal differential forms of a ρ -algebra

In this subsection we define the algebra of universal differential forms of the ρ -algebra A and we prove the universality of the construction. First we introduce the universal derivation of the order (α, β) , $\alpha, \beta \in G$.

Definition 3. Let M be a ρ -bimodule over A . The ρ -derivation $D : A \rightarrow M$ of the order (α, β) is *universal* if for ρ -derivation $D' : A \rightarrow N$ of the order $(\alpha, 0)$, there is a unique morphism $\Phi : M \rightarrow N$ of ρ -bimodules of degree β such that $D' = \Phi \circ D$.

Next we will construct an universal derivation over the ρ -algebra A .

Let $\alpha \in G$, $\mu : A \otimes A \rightarrow A$ be the map $\mu(x \otimes y) = \rho(\alpha, |y|)xy$ and $\Omega_\alpha^1 A = \ker(\mu)$.

We define the map

$$(2) \quad d : A \rightarrow \Omega_\alpha^1 A$$

by

$$(3) \quad dx = 1 \otimes x - \rho(\alpha, |x|)x \otimes 1,$$

for any $x \in A$.

Proposition 2. $\Omega_\alpha^1 A$ is a ρ -bimodule over A and the map $d : A \rightarrow \Omega_\alpha^1 A$ is an universal derivation of the order $(\alpha, 0)$.

Proof. We have to show that $a(b \otimes c) \in \Omega_\alpha^1 A$, $(b \otimes c)a \in \Omega_\alpha^1 A$ and $(a(b \otimes c))d = a((b \otimes c)d)$, for any $a, d \in A$ and $b \otimes c \in \Omega_\alpha^1 A$. Indeed

$$\mu(a(b \otimes c)) = \mu(ab \otimes c) = \rho(\alpha, |ab|)abc = \rho(\alpha, |a|)a(\rho(\alpha, |b|)bc) = 0.$$

The other relations are obvious.

It is easy to see that $\Omega_\alpha^1 A$ is the space $\{xdy : x, y \in A \text{ and } d \text{ from (3)}\}$.

Now we show that $d : A \rightarrow \Omega_\alpha^1 A$ is a ρ -derivation of the order $(\alpha, 0)$:

$$\begin{aligned} (da)b + \rho(\alpha, |a|)adb &= (1 \otimes a - \rho(\alpha, |a|)a \otimes 1)b + \rho(\alpha, |a|)a(1 \otimes b - \rho(\alpha, |b|)b \otimes 1) \\ &= 1 \otimes ab - \rho(\alpha, |a|)a \otimes b + \rho(\alpha, |a|)a \otimes b - \rho(\alpha, |a| + |b|)ab \otimes 1 \\ &= 1 \otimes (ab) - \rho(\alpha, |ab|)(ab) \otimes 1 = d(ab). \end{aligned}$$

Let M be a ρ -bimodule over A and $D:A \rightarrow M$ a ρ -derivation of the order (α, β) . We define the map $\Phi : \Omega_\alpha^1 A \rightarrow M$ in the following way:

$$\Phi(a \otimes b) = \rho(\alpha, |a|)aD(b),$$

for any $a \otimes b \in \Omega_\alpha^1 A$.

Now we show that Φ is a morphism of ρ -bimodules over A :

$$\begin{aligned} \Phi(a(b \otimes c)) &= \phi((ab) \otimes c) = \rho(\alpha, |ab|)abD(c) \\ &= \rho(\alpha, |a|)a[\rho(\alpha, |b|)bD(c)] = \rho(\alpha, |a|)a\Phi(b \otimes c). \end{aligned}$$

On the other hand

$$\begin{aligned} \Phi((a \otimes b)c) &= \Phi(a \otimes (bc)) = \rho(\alpha, |a|)aD(bc) \\ &= \rho(\alpha, |a|)a[D(b)c + \rho(\alpha, |b|)bD(c)] \\ &= \rho(\alpha, |a|)aD(b)c + \rho(\alpha, |b|)[\rho(\alpha, |a|)ab]D(c) \\ &= \rho(\alpha, |a|)aD(b)c + \rho(\alpha, |b|)\mu(a \otimes b)D(c) \\ &\stackrel{a \otimes b \in \ker(\mu)}{=} \rho(\alpha, |a|)aD(b)c = \Phi(a \otimes b)c. \end{aligned}$$

Finally we get

$$\begin{aligned} (\Phi \circ d)(a) &= \Phi(1 \otimes a - \rho(\alpha, |a|)a \otimes 1) \\ &= \rho(\alpha, |1|)D(a) - \rho(\alpha, |a|)\rho(\alpha, |a|)aD(1) = D(a) \end{aligned}$$

so we have proved that $\Phi \circ d = D$. □

Let

$$\Omega_\alpha^n A = \underbrace{\Omega_\alpha^1 A \otimes_A \dots \otimes_A \Omega_\alpha^1 A}_{n \text{ times}}, \quad \Omega_\alpha^0 A = A \quad \text{and} \quad \Omega_\alpha A = \bigoplus_{n \geq 0} \Omega_\alpha^n A.$$

Naturally $\Omega_\alpha A$ is a ρ -bimodule over A and a algebra with the multiplication $(\omega, \theta) \mapsto \omega \otimes_A \theta$, for any $\omega, \theta \in \Omega_\alpha A$.

Remark 1. The algebra $\Omega_\alpha A$ may be identified with the algebra generated by the algebra A and the derivations $da, a \in A$ which satisfies the following relations:

1. da is linear in a
2. the ρ -Leibniz rule: $d(ab) = d(a)b + \rho(\alpha, |a|)adb$
3. $d(1) = 0$.

$\Omega_\alpha^n A$ is the space of n -forms $a_0 da_1 \dots da_n$, $a_i \in A$ for any $0 \leq i \leq n$. $\Omega_\alpha^n A$ is a ρ -bimodule over A with the left multiplication

$$(4) \quad a(a_0 da_1 \dots da_n) = aa_0 da_1 \dots da_n,$$

and with the right multiplication given by:

$$\begin{aligned} (a_0 da_1 \dots da_n) a_{n+1} &= \sum_{i=1}^n (-1)^{n-i} \rho(\alpha, \sum_{j=i+1}^n |a_j|) (a_0 da_1 \dots d(a_i a_{i+1}) \dots da_{n+1}) \\ &\quad + (-1)^n \rho(\alpha, \sum_{i=1}^n |a_j|) a_0 a_1 da_2 \dots da_{n+1}. \end{aligned}$$

The multiplication in the algebra $\Omega_\alpha A$ is:

$$(a_0 da_1 \dots da_n)(a_{n+1} da_{n+2} \dots da_{m+n}) = ((a_0 da_1 \dots da_n) a_{n+1}) da_{n+2} \dots da_{m+n},$$

for any $a_i \in A$, $0 \leq i \leq n + m$, $n, m \in \mathbb{N}$.

We define the G -degree of the n -form $a_0 da_1 \dots da_n$ in the following way

$$|a_0 da_1 \dots da_n| = \sum_{i=0}^n |a_i|.$$

It is obvious that $|\omega_n \cdot \omega_m| = |\omega_n| + |\omega_m|$ for any homogeneous forms $\omega_n \in \Omega_\alpha^n A$ and $\omega_m \in \Omega_\alpha^m A$.

$\Omega_\alpha A$ is a $G' = \mathbb{Z} \times G$ -graded algebra with the G' degree of the n -form $a_0 da_1 \dots da_n$ defined by

$$|a_0 da_1 \dots da_n|' = \left(n, \sum_{i=0}^n |a_i| \right).$$

We may define the cocycle $\rho' : G' \times G' \rightarrow k$ on the algebra $\Omega_\alpha A$ thus:

$$(5) \quad \rho'(|a_0 da_1 \dots da_n|', |b_0 db_1 \dots db_m|') = (-1)^{nm} \rho \left(\sum_{i=0}^n |a_i|, \sum_{i=0}^m |b_i| \right)$$

and thus we have that $\Omega_\alpha A$ is a ρ' -algebra. It may be proved that the map $d : \Omega_\alpha A \rightarrow \Omega_\alpha A$ is a derivation of the order $((1, \alpha), (1, 0))$. Concluding we have the following result:

Theorem 1. $(\Omega_\alpha A, d)$ is a ρ -differential calculus over A .

Example 1. In the case when the group G is trivial then A is the usual associative algebra and $\Omega_\alpha A$ is the algebra of universal differential forms of A .

Example 2. If the group G is \mathbb{Z}_2 and the cocycle ρ is defined by $\rho(a, b) = (-1)^{ab}$ then A is a superalgebra. In the case when $\alpha = 1$ $\Omega_\alpha A$ is the superalgebra of universal differential forms of A from [11].

3. THE FROBENIUS THEOREM ON QUANTUM HYPERPLANE

In this section we give the Frobenius theorem on the N -dimensional quantum hyperplane S_N^q and we give the equations of any globally integrable distributions and parallel with respect to a connection ∇ . These results are valid for any ρ -differential calculus on S_N^q . In the first subsection we present two different ρ -differential calculi on S_N^q : the first one is the algebra of forms $\Omega(S_N^q)$ on S_N^q form [1] and the second one is the algebra of universal differential forms $\Omega_\alpha(S_N^q)$ on S_N^q . Both of these ρ -differential calculi on S_N^q are different by differential calculus on S_N^q of Wess and Zumino [15]. In the second subsection we present submanifolds on S_N^q , in the third subsection we review the basic notions about linear connections on S_N^q and finally we will give the main results of this paper.

3.1. Differential calculi on S_N^q

First we review the basic notions about the N -dimensional quantum hyperplane S_N^q . For more details see [1] and [3].

The N -dimensional quantum hyperplane S_N^q is the k -algebra generated by the unit element and N linearly independent elements x_1, \dots, x_N satisfying the relations: $x_i x_j = q x_j x_i$, $i < j$ for some fixed $q \in k$, $q \neq 0$.

S_N^q is a \mathbb{Z}^N -graded algebra

$$S_N^q = \bigoplus_{n_1, \dots, n_N}^{\infty} (S_N^q)_{n_1, \dots, n_N},$$

with $(S_N^q)_{n_1, \dots, n_N}$ the one-dimensional subspace spanned by products $x^{n_1} \dots x^{n_N}$. The \mathbb{Z}^N -degree of $x^{n_1} \dots x^{n_N}$ is $n = (n_1, \dots, n_N)$. The cocycle $\rho : \mathbb{Z}^N \times \mathbb{Z}^N \rightarrow k$ is

$$\rho(n, n') = q^{\sum_{j,k=1}^N n_j n'_k \alpha_{jk}},$$

with $\alpha_{jk} = 1$ for $j < k$, 0 for $j = k$ and -1 for $j > k$.

It may be proved that the N -dimensional quantum hyperplane S_N^q is a ρ -commutative algebra.

Remark that the space of ρ -derivations $\rho\text{-Der}(S_N^q)$ is a free S_N^q -module of rank N with $\partial/\partial x_1, \dots, \partial/\partial x_N$ as the basis, where $\partial/\partial x_i(x_j) = \delta_{ij}$.

Remark 2. Let $(\Omega S_N^q, d)$ a ρ -differential calculus on S_N^q , with $d : \Omega S_N^q \rightarrow \Omega S_N^q$ a ρ -derivation of the order $((1, \alpha), (1, 0))$, where $\alpha, 0 \in \mathbb{Z}^N$. It is easy to see that ΩS_N^q is generated by $\{x_1, \dots, x_N\}$ and there differentials $\{y_1 = dx_1, \dots, y_N = dx_N\}$ with some relations between them.

Next we give some examples of ρ -differential calculi on S_N^q .

3.1.1. The algebra of forms $\Omega(S_N^q)$ of S_N^q . $\Omega(S_N^q)$ ([1], [3]) is the algebra determined by the elements x_1, \dots, x_N and $y_1 = dx_1, \dots, y_N = dx_N$ with the relations

$$(6) \quad x_j x_k = q^{\alpha_{jk}} x_k x_j, \quad y_j y_k = -q^{\alpha_{jk}} y_k y_j, \quad x_j y_k = q^{\alpha_{jk}} y_k x_j.$$

3.1.2. The algebra of universal differential forms $\Omega_\alpha(A)$ of S_N^q . Next we will apply the construction of the algebra of the universal differential forms of a ρ -algebra from the remark 1 to the ρ -algebra S_N^q and, thus, we will give a new differential calculus on S_N^q denoted by $\Omega_\alpha(S_N^q)$.

Let $\alpha = (n_1, \dots, n_N)$ be an arbitrary element from \mathbb{Z}^N . $\Omega_\alpha(S_N^q)$ is the algebra generated by $a \in S_N^q$ and the symbols da , which satisfies the following relations:

1. da is linear in a .
2. the ρ -Leibniz rule: $d(ab) = (da)b + \rho(n, |a|)adb$.
3. $d(1) = 0$.

Next we present the structure of the algebra $\Omega_\alpha(S_N^q)$.

We use the following notations $y_i = dx_i$, for any $i \in \{1, \dots, N\}$. By an easy computation we get the following lemmas:

Lemma 1. $y_i x_j = \rho(\alpha + |x_i|, |x_j|) x_j y_i$, for any $i, j \in \{1, \dots, N\}$.

Lemma 2. $y_j y_i = \rho(\alpha, |x_i|) \rho(n + |x_i|, |x_j|) y_i y_j$, for any $i, j \in \{1, \dots, N\}$.

Lemma 3. $d(x_i^m) = m\rho^{m-1}(\alpha, |x_i|)x_i^{m-1}y_i$, for any $m \in \mathbb{N}$ and $i \in \{1, \dots, N\}$.

Putting together the previous lemmas we obtain the following theorem which gives the structures of the algebra $\Omega_\alpha(S_N^q)$:

Theorem 2. $\Omega_\alpha(S_N^q)$ is the algebra spanned by the elements x_i and y_i with $i \in \{1, \dots, N\}$ which satisfies the following relations:

1. $x_i x_j = \rho(|x_i|, |x_j|)x_j x_i$,
2. $y_i x_j = \rho(\alpha + |x_i|, |x_j|)x_j y_i$,
3. $y_j y_i = \rho(\alpha, |x_i|)\rho(\alpha + |x_i|, |x_j|)y_i y_j$, for any $i, j \in \{1, \dots, N\}$.

3.2. Submanifolds in S_N^q

In this section we use the definition of submanifolds algebra in noncommutative geometry from [11] to introduce submanifolds in the quantum hyperplane.

Let C be an ideal in S_N^q . We denote by $Q = S_N^q/C$ the quotient algebra and by $p : S_N^q \rightarrow Q$ the quotient map. We consider the following two Lie ρ -subalgebras of $\rho\text{-Der}S_N^q$:

$$G_C = \{X \in \rho\text{-Der}S_N^q/XC \subset C\}$$

and

$$G_A = \{X \in \rho\text{-Der}S_N^q/X(S_N^q) \subset C\}$$

We define the map $\pi : G_C \rightarrow \rho\text{-Der}S_N^q$ by $\pi(X)p(a) = p(Xa)$ for any $a \in S_N^q$ and $X \in G_C$.

Definition 4. The quotient algebra $Q = S_N^q/C$ is a *submanifold algebra* of S_N^q if the map π is surjective.

In this situation we obtain the following the short exact sequence of ρ -Lie algebras.

$$(7) \quad 0 \rightarrow G_A \rightarrow G_C \rightarrow \rho\text{-Der}Q \rightarrow 0$$

Let C be the ideal from S_N^q generated by the elements x_1, x_2, \dots, x_p with $p \in \{1, \dots, N\}$. Then the algebra Q generated by the elements x_{p+1}, \dots, x_N and we have that $A = C \oplus Q$.

Theorem 3. $G_C = (\rho\text{-Der}Q) \oplus G_A$.

Proof. Let X be a colour derivation from $\rho\text{-Der}S_N^q$, then

$$X = \sum_{i=1}^N X_i \frac{\partial}{\partial x_i} \quad \text{and} \quad X_i = X_i^C + X_i^Q$$

with $X_i^C \in C$ and $X_i^Q \in Q$ for any $i \in \{1, \dots, N\}$.

If $X \in G_C$ then $X(c) \in C$ for any $c \in C$ so

$$\begin{aligned} X(c) &= \sum_{i=1}^N X_i^C \frac{\partial c}{\partial x_i} + \sum_{i=1}^N X_i^Q \frac{\partial c}{\partial x_i} \\ &= \underbrace{\sum_{i=1}^N X_i^C \frac{\partial c}{\partial x_i}}_{\in C} + \sum_{i=1}^p X_i^Q \frac{\partial c}{\partial x_i} + \underbrace{\sum_{i=p+1}^N X_i^Q \frac{\partial c}{\partial x_i}}_{\in C} \in C \end{aligned}$$

Results that $X_i^Q = 0$ for any $i \in \{1, \dots, p\}$.

So any element $X = \sum_{i=1}^N X_i \partial/\partial x_i$ from G_C may be written is the following way

$$X = X^{G_A} + X^Q$$

where

$$X^{G_A} = \sum_{i=1}^N X_i^C \frac{\partial}{\partial x_i} \in G_A \quad \text{and} \quad X^Q = \sum_{i=1}^N X_i^Q \frac{\partial}{\partial x_i} \in \rho\text{-Der}Q$$

with $X_i^Q = 0$ for $i \in \{1, \dots, p\}$. □

Corollary 1. *Q is submanifolds algebra of S_N^q .*

3.3. Linear connections on S_N^q

In this subsection we use the definition of linear connections on ρ -algebras from the paper [3]. Let $(\Omega S_N^q, d)$ be a ρ -differential calculus on S_N^q and $n \in \mathbb{Z}^N$. A linear connection along the field $X = \sum_{i=1}^N X_i \frac{\partial}{\partial x_i}$ on the ρ -bimodule $\Omega^n S_N^q$ over S_N^q is a linear map

$$\nabla : \rho\text{-Der}(S_N^q) \rightarrow \text{End}(\Omega^n S_N^q)$$

of degree $|X|$ such that

$$\nabla(X)(a\omega) = \nabla_X(a\omega) = \rho(|X|, |\omega|)X(a)\omega + a\nabla_X\omega$$

for any $X \in \rho\text{-Der}S_N^q$, $a \in S_N^q$ and $\omega \in \Omega^n S_N^q$.

Using the structure of the free bimodule $\Omega^n S_N^q$ we deduce that any such connection ∇ is well defined by the connections coefficients $\Gamma_{i,i_1,\dots,i_n}^{j_1,\dots,j_n} \in S_N^q$ defined by

$$(8) \quad \nabla_{\frac{\partial}{\partial x_i}}(y_{i_1} \dots y_{i_n}) = \Gamma_{i,i_1,\dots,i_n}^{j_1,\dots,j_n} y_{j_1} \dots y_{j_n}.$$

Remark 3. The connection coefficients $\Gamma_{i,i_1,\dots,i_n}^{j_1,\dots,j_n}$ satisfy the some properties which depend on the choice of the ρ -differential calculus $(\Omega S_N^q, d)$.

Example 3. If $(\Omega S_N^q, d)$ is the algebra of forms $\Omega(S_N^q)$ of S_N^q we obtain that

$$(9) \quad \Gamma_{i, i_1, \dots, i_k, i_{k+1}, \dots, i_n}^{j_1, \dots, j_n} = -q^{\alpha_{i_k, i_{k+1}}} \Gamma_{i, i_1, \dots, i_{k+1}, i_k, \dots, i_n}^{j_1, \dots, j_n}$$

for any $i, i_1, \dots, i_n, j_1, \dots, j_n \in \{1, \dots, n\}$.

Example 4. If $(\Omega S_N^q, d)$ is the algebra of universal differential forms $\Omega_\alpha(S_N^q)$ of S_N^q using an easy computation we obtain that

$$(10) \quad \Gamma_{i, i_1, \dots, i_k, i_{k+1}, \dots, i_m}^{j_1, \dots, j_m} = \rho(\alpha, |y_{i_k}|) \rho(\alpha + |y_{i_k}|, |y_{i_{k+1}}|) \Gamma_{i, i_1, \dots, i_{k+1}, i_k, \dots, i_m}^{j_1, \dots, j_m}.$$

3.4. Distributions in S_N^q

Next we introduce distributions on S_N^q . Let $(\Omega S_N^q, d)$ be a ρ -differential calculus on S_N^q .

Definition 5. A *distribution* \mathcal{D} on S_N^q is a S_N^q -subbimodule of $\Omega^1 S_N^q$. The distribution \mathcal{D} is *globally integrable* if there is a subset B of S_N^q such that \mathcal{D} is the subspace generated by $S_N^q d(B)$ and by $d(B) S_N^q$.

Definition 6. We say that the distribution \mathcal{D} is *parallel* with respect to the connection $\nabla : \rho\text{-Der} S_N^q \rightarrow \text{End}(\Omega^1 S_N^q)$ if

$$\nabla_X(m) = 0, \text{ for any } X \in \rho\text{-Der} S_N^q \text{ and for any } m \in \mathcal{D}.$$

Using the structure of ΩS_N^q we obtain the following structure theorem of globally integrable distributions.

Theorem 4. *Any globally integrable distributions \mathcal{D} determined by $S_N^q d(B)$ and $d(B) S_N^q$ where B is the subset $\{x_1, \dots, x_p\}$. In this situation we say that the distribution \mathcal{D} has the dimension p .*

3.4.1. The Frobenius theorem for quantum hyperplane. In this section we will give a Frobenius theorem for N -dimensional quantum hyperplane which is obvious from the previous results.

Theorem 5. *The Frobenius theorem for quantum hyperplane. Any globally integrable distribution from S_N^q is given by a maximal submanifold algebra of S_N^q and conversely any submanifold algebra of S_N^q give a globally integrable distribution with the same dimension.*

Proof. Let \mathcal{D} be a global integrable distribution from S_N^q . Then from the theorem 10, \mathcal{D} is given by $S_N^q dQ$, where Q is a subset with p elements from $\{x_1, \dots, x_N\}$. If we denote by C the ideal of S_N^q generated by $\{x_1, \dots, x_N\} \setminus Q$ and using the Corollary 9 we obtain that $Q = S_N^q / C$ is a submanifold algebra of S_N^q of the dimension p .

Conversely, if Q is a submanifold algebra of the dimension p of S_N^q results that there is an ideal C of the dimension $N - p$ of S_N^q such that $Q = S_N^q / C$. If we denote by $\{x_1, \dots, x_{N-p}\}$ is the subset of $\{x_1, \dots, x_N\}$ which generates the ideal C then the set $Q = \{x_{N-p+1}, \dots, x_N\}$ generates a distribution of the dimension p of S_N^q . \square

We may find the equations of an globally integrable distributions and parallel with respect to a connection ∇ .

Theorem 6. *Any globally integrable and parallel distribution \mathcal{D} with respect to a connection $\nabla : \rho\text{-Der}S_N^q \rightarrow \text{End}(\Omega^1 S_N^q)$ of dimension p is given by the following equations:*

$$(11) \quad \Gamma_{i,j}^k = 0$$

for a subset I of $\{1, \dots, N\}$ with p elements and for any $i \in \{1, \dots, N\}$, $j \in I$, $k \in \{1, \dots, N\} \setminus I$.

1. Bongaarts P. J. M. and Pijls H. G. J., *Almost commutative algebra and differential calculus on the quantum hyperplane.* J. Math. Phys. **35**(2) (1994), 959-970.
2. Cap A., Kriegl A., Michor P. W. and Vanžura J., *The Frölicher-Nijenhuis bracket in noncommutative differential geometry.* Acta Math. Univ. Comenianiae **62** (1993), 17-49.
3. Ciupală C., *Linear connections on almost commutative algebras.* Acta Math. Univ. Comenianiae **72**(2) (2003), 197-207.
4. ———, *ρ -differential calculi and linear connections on matrix algebra.* Int. J. Geom. Methods in Mod. Phys. **1** (2004), 847-863.
5. ———, *Fields and forms on ρ -algebras.* Proc. Indian Acad. Sci. -Math. Sciences. **115** (2005), 57-65.

6. ———, *Linear connections on quaternionic algebra*, Manuscript in preparation.
7. Connes A., *Non-commutative Geometry*. Academic Press, 1994.
8. Dubois-Violette M. and Michor P. W., *Connections on central bimodules*, J. Geom. Phys. **20** (1996), 218–232.
9. Dubois-Violette M., *Lectures on graded differential algebras and noncommutative geometry*. Vienne, Preprint, E.S.I. **842** (2000).
10. Jadczyk A., Kastler D., *Graded Lie-Cartan pairs. II. The fermionic differential calculus*. Ann. of Physics **179** (1987), 169–200.
11. Kastler D., *Cyclic Cohomology within the differential envelope*. Hermann, Paris 1988.
12. Lychagin V., *Colour calculus and colour quantizations*. Acta Appl. Math. **41** (1995), 193–226.
13. Mourad J., *Linear connections in noncommutative geometry*. Class. Quantum Grav. **12** (1995) 965.
14. Masson T., *Submanifolds and quotient manifolds in noncommutative geometry*. Lett. Math. Phys. **37** (1996), 2484–2497.
15. Wess J. and Zumino B., *Differential calculus on quantum hyperplane*. Nuclear Physics **18B** (1990), 303–312.
16. Woronowicz S. L., *Differential calculus on compact matrix pseudogroups (quantum groups)*. Comm. Math. Phys. **112**(1) (1989), 125–170.

C. Ciupală, The "Andrei Şaguna" National College, Andrei Şaguna str. No. 1, Braşov, Romania,
e-mail: cciupala@yahoo.com