

DIFFERENTIAL SANDWICH THEOREMS FOR SOME SUBCLASSES OF ANALYTIC FUNCTIONS INVOLVING A LINEAR OPERATOR

T. N. SHAMMUGAM, C. RAMACHANDRAN, M. DARUS AND S. SIVASUBRAMANIAN

ABSTRACT. By making use of the familiar Carlson–Shaffer operator, the authors derive some subordination and superordination results for certain normalized analytic functions in the open unit disk. Relevant connections of the results, which are presented in this paper, with various other known results are also pointed out.

1. INTRODUCTION

Let \mathcal{H} be the class of functions *analytic* in the *open* unit disk

$$\Delta := \{z : |z| < 1\}.$$

Let $\mathcal{H}[a, n]$ be the subclass of \mathcal{H} consisting of functions of the form

$$f(z) = a + a_n z^n + a_{n+1} z^{n+1} + \dots$$

Let

$$\mathcal{A}_m := \{f \in \mathcal{H}, f(z) = z + a_{m+1} z^{m+1} + a_{m+2} z^{m+2} + \dots\}$$

and let $\mathcal{A} := \mathcal{A}_1$. With a view to recalling the principle of subordination between analytic functions, let the functions f and g be analytic in Δ . Then we say that the function f is *subordinate* to g if there exists a Schwarz

Received August 12, 2006.

2000 *Mathematics Subject Classification.* Primary 30C80; Secondary 30C45.

Key words and phrases. differential subordinations, differential superordinations, sominant, subordinant.

function ω , analytic in Δ with

$$\omega(0) = 0 \quad \text{and} \quad |\omega(z)| < 1 \quad (z \in \Delta),$$

such that

$$f(z) = g(\omega(z)) \quad (z \in \Delta).$$

We denote this subordination by

$$f \prec g \quad \text{or} \quad f(z) \prec g(z).$$

In particular, if the function g is univalent in Δ , the above subordination is equivalent to

$$f(0) = g(0) \quad \text{and} \quad f(\Delta) \subset g(\Delta).$$

Let $p, h \in \mathcal{H}$ and let $\phi(r, s, t; z) : \mathbb{C}^3 \times \Delta \rightarrow \mathbb{C}$. If p and $\phi(p(z), zp'(z), z^2p''(z); z)$ are univalent and if p satisfies the second order superordination

$$(1.1) \quad h(z) \prec \phi(p(z), zp'(z), z^2p''(z); z),$$

then p is a solution of the differential superordination (1.1). (If f is subordinate to F , then F is called to be superordinate to f .) An analytic function q is called a *subordinant* if $q \prec p$ for all p satisfying (1.1). An univalent subordinant \tilde{q} that satisfies $q \prec \tilde{q}$ for all subordinants q of (1.1) is said to be the best subordinant. Recently Miller and Mocanu [6] obtained conditions on h , q and ϕ for which the following implication holds:

$$h(z) \prec \phi(p(z), zp'(z), z^2p''(z); z) \Rightarrow q(z) \prec p(z).$$

with the results of Miller and Mocanu [6], Bulboacă [3] investigated certain classes of first order differential subordinations as well as superordination-preserving integral operators [2]. Ali et al. [1] used the results obtained by Bulboacă [3] and gave sufficient conditions for certain normalized analytic functions f to satisfy

$$q_1(z) \prec \frac{zf'(z)}{f(z)} \prec q_2(z)$$

where q_1 and q_2 are given univalent functions in Δ with $q_1(0) = 1$ and $q_2(0) = 1$. Shanmugam et al. [7] obtained sufficient conditions for a normalized analytic functions f to satisfy

$$q_1(z) \prec \frac{f(z)}{zf'(z)} \prec q_2(z)$$

and

$$q_1(z) \prec \frac{z^2 f'(z)}{(f(z))^2} \prec q_2(z).$$

where q_1 and q_2 are given univalent functions in Δ with $q_1(0) = 1$ and $q_2(0) = 1$. Recently, the first author combined with the third and fourth authors of this paper obtained sufficient conditions for certain normalized analytic functions f to satisfy

$$q_1(z) \prec \frac{z}{L(a, c)f(z)} \prec q_2(z)$$

where q_1 and q_2 are given univalent functions with $q_1(0) = 1$ and $q_2(0) = 1$ (see [8] for details; also see [9]). A detailed investigation of starlike functions of complex order and convex functions of complex order using Briot-Bouquet differential subordination technique has been studied very recently by Srivastava and Lashin [10].

Let the function $\varphi(a, c; z)$ be given by

$$\varphi(a, c; z) := \sum_{n=0}^{\infty} \frac{(a)_n}{(c)_n} z^{n+1} \quad (c \neq 0, -1, -2, \dots; z \in \Delta),$$

where $(x)_n$ is the *Pochhammer symbol* defined by

$$(x)_n := \begin{cases} 1, & n = 0; \\ x(x+1)(x+2)\dots(x+n-1), & n \in \mathbb{N} := \{1, 2, 3, \dots\}. \end{cases}$$

Corresponding to the function $\varphi(a, c; z)$, Carlson and Shaffer [4] introduced a linear operator $L(a, c)$, which is defined by the following Hadamard product (or convolution):

$$L(a, c)f(z) := \varphi(a, c; z) * f(z) = \sum_{n=0}^{\infty} \frac{(a)_n}{(c)_n} a_n z^{n+1}.$$

We note that

$$L(a, a)f(z) = f(z), \quad L(2, 1)f(z) = zf'(z), \quad L(\delta + 1, 1)f(z) = D^\delta f(z),$$

where $D^\delta f$ is the *Ruscheweyh derivative* of f .

The main object of the present sequel to the aforementioned works is to apply a method based on the differential subordination in order to derive several subordination results involving the Carlson Shaffer Operator. Furthermore, we obtain the previous results of Srivastava and Lashin [10] as special cases of some of the results presented here.

2. PRELIMINARIES

In order to prove our subordination and superordination results, we make use of the following known results.

Definition 1. [6, Definition 2, p. 817] Denote by Q the set of all functions f that are analytic and injective on $\overline{\Delta} - E(f)$, where

$$E(f) = \{\zeta \in \partial\Delta : \lim_{z \rightarrow \zeta} f(z) = \infty\},$$

and are such that $f'(\zeta) \neq 0$ for $\zeta \in \partial\Delta - E(f)$.

Theorem 1. [5, Theorem 3.4h, p. 132] *Let the function q be univalent in the open unit disk Δ and θ and ϕ be analytic in a domain D containing $q(\Delta)$ with $\phi(w) \neq 0$ when $w \in q(\Delta)$. Set $Q(z) = zq'(z)\phi(q(z))$, $h(z) = \theta(q(z)) + Q(z)$. Suppose that*

1. Q is starlike univalent in Δ , and

$$2. \Re \left(\frac{zh'(z)}{Q(z)} \right) > 0 \text{ for } z \in \Delta.$$

If

$$\theta(p(z)) + zp'(z)\phi(p(z)) \prec \theta(q(z)) + zq'(z)\phi(q(z)),$$

then $p(z) \prec q(z)$ and q is the best dominant.

Theorem 2. [3] Let the function q be univalent in the open unit disk Δ and ϑ and φ be analytic in a domain D containing $q(\Delta)$. Suppose that

1. $\Re \frac{\vartheta'(q(z))}{\varphi(q(z))} > 0$ for $z \in \Delta$,
2. $zq'(z)\varphi(q(z))$ is starlike univalent in Δ .

If $p \in \mathcal{H}[q(0), 1] \cap Q$, with $p(\Delta) \subseteq D$, and $\vartheta(p(z)) + zp'(z)\varphi(p(z))$ is univalent in Δ , and

$$\vartheta(q(z)) + zq'(z)\varphi(q(z)) \prec \vartheta(p(z)) + zp'(z)\varphi(p(z)),$$

then $q(z) \prec p(z)$ and q is the best subdominant.

3. SUBORDINATION AND SUPERORDINATION FOR ANALYTIC FUNCTIONS

We begin by proving involving differential subordination between analytic functions.

Theorem 3. Let $\left(\frac{L(a+1, c)f(z)}{z} \right)^\mu \in \mathcal{H}$ and let the function $q(z)$ be analytic and univalent in Δ such that $q(z) \neq 0, (z \in \Delta)$. Suppose that $\frac{zq'(z)}{q(z)}$ is starlike univalent in Δ . Let

$$(3.1) \quad \Re \left\{ 1 + \frac{\xi}{\beta} q(z) + \frac{2\delta}{\beta} (q(z))^2 - \frac{zq'(z)}{q(z)} + \frac{zq''(z)}{q'(z)} \right\} > 0$$

$$(\alpha, \delta, \xi, \beta \in \mathbb{C}; \beta \neq 0)$$

and

$$(3.2) \quad \begin{aligned} \Psi(a, c, \mu, \xi, \beta, \delta, f)(z) := & \alpha + \xi \left[\frac{L(a+1, c)f(z)}{z} \right]^\mu + \delta \left[\frac{L(a+1, c)f(z)}{z} \right]^{2\mu} \\ & + \beta\mu(a+2) \left[\frac{L(a+2, c)f(z)}{L(a+1, c)f(z)} - 1 \right]. \end{aligned}$$

If q satisfies the following subordination:

$$\Psi(a, c, \mu, \xi, \beta, \delta, f)(z) \prec \alpha + \xi q(z) + \delta(q(z))^2 + \beta \frac{zq'(z)}{q(z)}$$

$$(\alpha, \delta, \xi, \beta, \mu \in \mathbb{C}; \mu \neq 0; \beta \neq 0),$$

then

$$(3.3) \quad \left(\frac{L(a+1, c)f(z)}{z} \right)^\mu \prec q(z) \quad (\mu \in \mathbb{C}; \mu \neq 0)$$

and q is the best dominant.

Proof. Let the function p be defined by

$$p(z) := \left(\frac{L(a+1, c)f(z)}{z} \right)^\mu \quad (z \in \Delta; z \neq 0; f \in \mathcal{A}),$$

so that, by a straightforward computation, we have

$$\frac{zp'(z)}{p(z)} = \mu \left[\frac{z(L(a+1, c)f(z))'}{L(a+1, c)f(z)} - 1 \right].$$

By using the identity:

$$z(L(a, c)f(z))' = (1 + a)L(a + 1, c)f(z) - aL(a, c)f(z),$$

we obtain

$$\frac{zp'(z)}{p(z)} = \mu \left[(a + 2) \frac{L(a + 2, c)f(z)}{L(a + 1, c)f(z)} - (a + 2) \right].$$

By setting

$$\theta(\omega) := \alpha + \xi\omega + \delta\omega^2 \quad \text{and} \quad \phi(\omega) := \frac{\beta}{\omega},$$

it can be easily observed that θ is analytic in \mathbb{C} , ϕ is analytic in $\mathbb{C} \setminus \{0\}$ and that

$$\phi(\omega) \neq 0 \quad (\omega \in \mathbb{C} \setminus \{0\}).$$

Also, by letting

$$Q(z) = zq'(z)\phi(q(z)) = \beta \frac{zq'(z)}{q(z)}$$

and

$$h(z) = \theta(q(z)) + Q(z) = \alpha + \xi q(z) + \delta(q(z))^2 + \beta \frac{zq'(z)}{q(z)},$$

we find that $Q(z)$ is starlike univalent in Δ and that

$$\Re \left(\frac{zh'(z)}{Q(z)} \right) = \Re \left\{ 1 + \frac{\xi}{\beta} q(z) + \frac{2\delta}{\beta} (q(z))^2 - \frac{zq'(z)}{q(z)} + \frac{zq''(z)}{q'(z)} \right\} > 0,$$

$$(\alpha, \delta, \xi, \beta \in \mathbb{C}; \beta \neq 0).$$

The assertion (3.3) of Theorem 3 now follows by an application of Theorem 1. □

For the choices $q(z) = \frac{1 + Az}{1 + Bz}$, $-1 \leq B < A \leq 1$ and $q(z) = \left(\frac{1+z}{1-z}\right)^\gamma$, $0 < \gamma \leq 1$, in Theorem 3, we get the following results (Corollaries 1 and 2 below).

Corollary 1. Assume that (3.1) holds. If $f \in \mathcal{A}$, and

$$\Psi(a, c, \mu, \xi, \beta, \delta, f)(z) \prec \alpha + \xi \frac{1 + Az}{1 + Bz} + \delta \left(\frac{1 + Az}{1 + Bz}\right)^2 + \frac{\beta(A - B)z}{(1 + Az)(1 + Bz)}$$

$$(z \in \Delta; \alpha, \delta, \xi, \beta, \mu \in \mathbb{C}; \mu \neq 0; \beta \neq 0),$$

where $\Psi(a, c, \mu, \xi, \beta, \delta, f)$ is as defined in (3.2), then

$$\left(\frac{L(a+1, c)f(z)}{z}\right)^\mu \prec \frac{1 + Az}{1 + Bz} \quad (\mu \in \mathbb{C}; \mu \neq 0)$$

and $\frac{1 + Az}{1 + Bz}$ is the best dominant.

Corollary 2. Assume that (3.1) holds. If $f \in \mathcal{A}$, and

$$\Psi(a, c, \mu, \xi, \beta, \delta, f)(z) \prec \alpha + \xi \left(\frac{1 + z}{1 - z}\right)^\gamma + \delta \left(\frac{1 + z}{1 - z}\right)^{2\gamma} + \frac{2\beta\gamma z}{(1 - z)^2}$$

$$(\alpha, \delta, \xi, \beta, \mu \in \mathbb{C}; \mu \neq 0; \beta \neq 0)$$

where $\Psi(a, c, \mu, \xi, \beta, \delta, f)(z)$ is as defined in (3.2), then

$$\left(\frac{L(a+1, c)f(z)}{z}\right)^\mu \prec \left(\frac{1 + z}{1 - z}\right)^\gamma \quad (\mu \in \mathbb{C}; \mu \neq 0)$$

and $\left(\frac{1 + z}{1 - z}\right)^\gamma$ is the best dominant.

For a special case $q(z) = e^{\mu Az}$, with $|\mu A| < \pi$, Theorem 3 readily yields the following.

Corollary 3. *Assume that (3.1) holds. If $f \in \mathcal{A}$, and*

$$\Psi(a, c, \mu, \xi, \beta, \delta, f)(z) \prec \alpha + \xi e^{\mu Az} + \delta e^{2\mu Az} + \beta A \mu z$$

$$(\alpha, \delta, \xi, \beta, \mu \in \mathbb{C}; \mu \neq 0; \beta \neq 0)$$

where $\Psi(a, c, \mu, \xi, \beta, \delta, f)(z)$ is as defined in (3.2), then

$$\left(\frac{L(a+1, c)f(z)}{z} \right)^\mu \prec e^{\mu Az} \quad (\mu \in \mathbb{C}, \mu \neq 0)$$

and $e^{\mu Az}$ is the best dominant.

For a special case when $q(z) = \frac{1}{(1-z)^{2b}}$ ($b \in \mathbb{C} \setminus \{0\}$), $a = c = 1$, $\delta = \xi = 0$, $\mu = \alpha = 1$ and $\beta = \frac{1}{b}$, Theorem 3 reduces at once to the following known result obtained by Srivastava and Lashin [10].

Corollary 4. *Let b be a non zero complex number. If $f \in \mathcal{A}$, and*

$$1 + \frac{1}{b} \frac{zf''(z)}{f'(z)} \prec \frac{1+z}{1-z},$$

then

$$f'(z) \prec \frac{1}{(1-z)^{2b}}$$

and $\frac{1}{(1-z)^{2b}}$ is the best dominant.

Next, by appealing to Theorem 2 of the preceding section, we prove Theorem 4 below.

Theorem 4. Let q be analytic and univalent in Δ such that $q(z) \neq 0$ and $\frac{zq'(z)}{q(z)}$ be starlike univalent in Δ . Further, let us assume that

$$(3.4) \quad \Re \left[\frac{2\delta}{\beta} (q(z))^2 + \frac{\xi}{\beta} q(z) \right] > 0, \quad (\delta, \xi, \beta \in \mathbb{C}; \beta \neq 0).$$

If $f \in \mathcal{A}$,

$$0 \neq \left(\frac{L(a+1, c)f(z)}{z} \right)^\mu \in \mathcal{H}[q(0), 1] \cap \mathcal{Q},$$

and $\Psi(a, c, \mu, \xi, \beta, \delta, f)$ is univalent in Δ , then

$$\alpha + \xi q(z) + \delta (q(z))^2 + \beta \frac{zq'(z)}{q(z)} \prec \Psi(a, c, \mu, \xi, \beta, \delta, f)$$

$$(z \in \Delta; \alpha, \delta, \xi, \beta, \mu \in \mathbb{C}; \mu \neq 0; \beta \neq 0),$$

implies

$$(3.5) \quad q(z) \prec \left(\frac{L(a+1, c)f(z)}{z} \right)^\mu \quad (\mu \in \mathbb{C}; \mu \neq 0)$$

and q is the best subordinator where $\Psi(a, c, \mu, \xi, \beta, \delta, f)(z)$ is as defined in (3.2).

Proof. By setting

$$\vartheta(w) := \alpha + \xi w + \delta w^2 \quad \text{and} \quad \varphi(w) := \beta \frac{1}{w},$$

it is easily observed that ϑ is analytic in \mathbb{C} . Also, φ is analytic in $\mathbb{C} \setminus \{0\}$ and that

$$\varphi(w) \neq 0, \quad (w \in \mathbb{C} \setminus \{0\}).$$

Since q is convex (univalent) function it follows that,

$$\Re \frac{\vartheta'(q(z))}{\varphi(q(z))} = \Re \left[\frac{2\delta}{\beta} (q(z))^2 + \frac{\xi}{\beta} q(z) \right] > 0,$$

$$(\delta, \xi, \beta \in \mathbb{C}; \beta \neq 0).$$

The assertion (3.5) of Theorem 4 follows by an application of Theorem 2. □

We remark here that Theorem 4 can easily be restated, for different choices of the function q . Combining Theorem 3 and Theorem 4, we get the following sandwich theorem.

Theorem 5. *Let q_1 and q_2 be univalent in Δ such that $q_1(z) \neq 0$ and $q_2(z) \neq 0$, ($z \in \Delta$) with $\frac{zq_1'(z)}{q_1(z)}$ and $\frac{zq_1'(z)}{q_1(z)}$ being starlike univalent. Suppose that q_1 satisfies (3.4) and q_2 satisfies (3.1). If $f \in \mathcal{A}$,*

$$\left(\frac{L(a+1, c)f(z)}{z} \right)^\mu \in \mathcal{H}[q(0), 1] \cap \mathcal{Q} \quad \text{and} \quad \Psi(a, c, \mu, \xi, \beta, \delta, f)(z)$$

is univalent in Δ , then

$$\alpha + \xi q_1(z) + \delta (q_1(z))^2 + \beta \frac{zq_1'(z)}{q_1(z)} \prec \Psi(a, c, \mu, \xi, \beta, \delta, f)(z)$$

$$\prec \alpha + \xi q_2(z) + \delta (q_2(z))^2 + \beta \frac{zq_2'(z)}{q_2(z)}$$

$$(\alpha, \delta, \xi, \beta, \mu \in \mathbb{C}; \mu \neq 0; \beta \neq 0),$$

implies

$$q_1(z) \prec \left(\frac{L(a+1, c)f(z)}{z} \right)^\mu \prec q_2(z) \quad (\mu \in \mathbb{C}; \mu \neq 0)$$

and q_1 and q_2 are respectively the best subordinant and the best dominant.

Corollary 5. Let q_1 and q_2 be univalent in Δ such that $q_1(z) \neq 0$ and $q_2(z) \neq 0$ ($z \in \Delta$) with $\frac{zq_1'(z)}{q_1(z)}$ and $\frac{zq_1'(z)}{q_1(z)}$ being starlike univalent. Suppose q_1 satisfies (3.4) and q_2 satisfies (3.1). If $f \in \mathcal{A}$, $(f')^\mu \in \mathcal{H}[q(0), 1] \cap \mathcal{Q}$ and let

$$\Psi_1(\mu, \xi, \beta, \delta, f) := \alpha + \xi [f'(z)]^\mu + \delta [f'(z)]^{2\mu} + \frac{3}{2}\beta\mu \frac{zf''(z)}{f'(z)}$$

is univalent in Δ , then

$$\begin{aligned} \alpha + \xi q_1(z) + \delta (q_1(z))^2 + \beta \frac{zq_1'(z)}{q_1(z)} &< \Psi_1(\mu, \xi, \beta, \delta, f) \\ &< \alpha + \xi q_2(z) + \delta (q_2(z))^2 + \beta \frac{zq_2'(z)}{q_2(z)} \end{aligned}$$

$$(\alpha, \delta, \xi, \beta, \mu \in \mathbb{C}; \mu \neq 0; \beta \neq 0),$$

implies

$$q_1(z) < (f')^\mu < q_2(z) \quad (\mu \in \mathbb{C}; \mu \neq 0)$$

and q_1 and q_2 are respectively the best subordinant and the best dominant.

Acknowledgment. The work presented here was partially supported by SAGA Grant: STGL-012-2006, Academy of Sciences, Malaysia. The authors also would like to thank the referee for critical comments and suggestions to improve the content of the paper.

1. Ali R. M., Ravichandran V., Hussain Khan M. and Subramanian K. G., *Differential sandwich theorems for certain analytic functions*, Far East J. Math. Sci., **15**(1) (2005), 87–94.
2. Bulboacă T., *A class of superordination-preserving integral operators*, Indag. Math. New Ser., **13**(3) (2002), 301–311.
3. ———, *Classes of first-order differential subordinations*, Demonstr. Math., **35**(2) (2002), 287–292.
4. Carlson B. C. and Shaffer D. B., *Starlike and prestarlike hypergeometric functions*, SIAM J. Math. Anal., **15** (1984), 737–745.
5. Miller S. S. and Mocanu P. T., *Differential Subordinations: Theory and Applications*, Pure and Applied Mathematics No. 225, Marcel Dekker, New York, (2000).
6. ———, *Subordinants of differential superordinations*, Complex Variables, **48**(10) (2003), 815–826.
7. Shanmugam T. N., Ravichandran V. and Sivasubramanian S., *Differential Sandwich Theorems for some subclasses of Analytic Functions*, Austral. J. Math. Anal. Appl., 3(1) (2006), Article 8, 11 pp (electronic).
8. Shanmugam T. N., Sivasubramanian S. and Darus M., *On certain subclasses of functions involving a linear Operator*, Far East J. Math. Sci., **23**(3), (2006), 329–339.
9. Shanmugam T. N., Sivasubramanian S. and Srivastava H. M., *Differential sandwich theorems for certain subclasses of analytic functions involving multiplier transformations*, Int. Transforms Spec. Functions., **17**(12) (2006), 889–899.
10. Srivastava H. M., and Lashin A. Y., *Some applications of the Briot-Bouquet differential subordination*, J. Inequal. Pure Appl. Math., **6**(2) (2005), Article 41, 7 pp.

T. N. Shammugam, Department of Information Technology, Salalah College of Technology, P.O. Box 608 Salalah, Sultanate of Oman,
e-mail: drtns2001@yahoo.com

C. Ramachandran, Department of Mathematics, Easwari Engineering College, Ramapuram, Chennai-600 089, India,
e-mail: crjsp2004@yahoo.com

M. Darus, School Of Mathematical Sciences, Faculty Of Sciences And Technology UKM, BANGI 43600, Malaysia,
e-mail: maslina@pkrisc.cc.ukm.my

S. Sivasubramanian, Department of Mathematics, Easwari Engineering College, Ramapuram, Chennai-600 089, India,
e-mail: sivasaisastha@rediffmail.com