# ON WEAKLY REVERSIBLE RINGS

ZHAO LIANG AND YANG GANG

ABSTRACT. Let R be a ring. We introduce weakly reversible rings, which are a generalization of reversible rings, and investigate their properties. Moreover, we show that a ring R is weakly reversible if and only if for any n, the n-by-n upper triangular matrix ring  $T_n(R)$  is weakly reversible. Also some kinds of examples needed in the process are given.

# 1. INTRODUCTION

Throughout this paper, all rings are associative with identity. According to Cohn [1], a ring R is called reversible if ab = 0 implies ba = 0 for  $a, b \in R$ . Anderson-Camillo [2], observing the rings whose zero products commute, used the term  $ZC_2$  for what is called reversible; while Krempa-Niewieczerzal [3] took the term  $C_0$  for it. A generalization of reversible rings is investigated in this paper. We call a ring R weakly reversible if ab = 0 implies that Rbra is a nil left ideal of R for all  $a, b, r \in R$ . Clearly semicommutative rings are weakly reversible. It is well known that semicommutative does not imply reversible (e.g. see [9, Examples 3.9 and 3.11]), examples are given to show that weakly reversible rings are not necessarily semicommutative. It is shown that a ring R is weakly reversible if and only if for any n, the n-by-n upper triangular matrix  $T_n(R)$  is a weakly reversible ring.

# 2. Weakly reversible rings

**Definition.** A ring R is called weakly reversible if for all  $a, b, r \in R$  such that ab = 0, Rbra is a nil left ideal of R (equivalently, braR is nil right ideal of R).

A ring R is called semicommutative if for all  $a, b \in R$ , ab = 0 implies aRb = 0. This is equivalent to the definition that any left(right) annihilator over R is an ideal of R [6, Lemma 1.1].

Since every reversible ring is semicommutative, clearly every reversible ring is weakly reversible. In the following we will see the converse is not true. Note the class of weakly reversible rings is closed under subrings and finite direct products.

Received December 23, 2005.

<sup>2000</sup> Mathematics Subject Classification. Primary 16S50, 16U80.

Key words and phrases. Reversible rings, weakly reversible rings, semicommutative rings.

**Proposition 2.1.** Let R be a ring and I an ideal of R such that R/I is weakly reversible. If  $I \subseteq nil(R)$ , then R is weakly reversible.

*Proof.* Let  $a, b \in R$  and ab = 0, then  $\bar{a}\bar{b} = \bar{0}$ . Thus $(\bar{r_1}\bar{b}\bar{r_2}\bar{a}\bar{r_3})^n = \bar{0}$  for some positive integer n, where  $\bar{r_1}, \bar{r_2}, \bar{r_3} \in R/I$ . Hence  $(r_1br_2ar_3)^n \in I \subseteq \operatorname{nil}(R)$ . This means that R is weakly reversible.

**Theorem 2.2.** Suppose S and T are rings, and M is an (S,T)-bimodule. Let

$$R = \left(\begin{array}{cc} S & M \\ 0 & T \end{array}\right).$$

Then R is weakly reversible if and only if S and T are weakly reversible.

*Proof.* Note that any subring of a weakly reversible ring is weakly reversible, so if R is weakly reversible then S and T are weakly reversible.

Conversely, suppose S and T are weakly reversible. Put

$$I = \left(\begin{array}{cc} 0 & M \\ 0 & 0 \end{array}\right) \triangleleft R.$$

The ring  $R/I \cong S \times T$  is weakly reversible, and now Proposition 2.1 implies that R is weakly reversible.

The following proposition follows immediately by induction on n.

**Proposition 2.3.** A ring R is a weakly reversible ring if and only if, for any n, the n-by-n upper triangular matrix ring  $T_n(R)$  is a weakly reversible ring.

Given a ring R and a bimodule  ${}_{R}M_{R}$ , the trivial extension of R by M is the ring  $T(R, M) = R \bigoplus M$  with the usual addition and the following multiplication

$$(r_1, m_1)(r_2, m_2) = (r_1r_2, r_1m_2 + m_1r_2).$$

This is isomorphic to the ring of all matrix  $\begin{pmatrix} r & m \\ 0 & r \end{pmatrix}$ , where  $r \in R, m \in M$  and the usual matrix operations are used.

**Corollary 2.4.** A ring R is a weakly reversible ring if and only if its trivial extension is a weakly reversible ring.

**Corollary 2.5.** Let R be a ring, then R is a weakly reversible ring if and only if for any  $n \in N$ ,  $R[x]/(x^n)$  is a weakly reversible ring, where  $(x^n)$  is the ideal of R[x] generated by  $x^n$ .

Now we can give examples of weakly reversible rings which are not reversible. As we know, reversible rings are both semicommutative [6, Lemma 1.4] and weakly reversible by definition. So we may conjecture that weakly reversible rings may be semicommutative. But the following example eliminates the possibility.

190

**Example 2.6.** Let S be a weakly reversible ring. Then

$$T = \left\{ \left( \begin{array}{ccc} a_{11} & a_{12} & a_{13} \\ 0 & a_{22} & a_{23} \\ 0 & 0 & a_{33} \end{array} \right) \, \middle| \, a_{ij} \in S \right\}$$

is a weakly reversible ring by Proposition 2.3. Note that

$$\left(\begin{array}{ccc} 0 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{array}\right) \left(\begin{array}{ccc} 0 & 1 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{array}\right) = 0$$

But we have

$$\left(\begin{array}{ccc} 0 & 1 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{array}\right) \left(\begin{array}{ccc} 0 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{array}\right) = \left(\begin{array}{ccc} 0 & 0 & 1 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{array}\right) \neq 0.$$

So T is not reversible. In fact, T is not semiccomutative by the argument from the last sentence of [9, Example 3.17] (with n=3).

Also let S be a weakly reversible ring. Then the ring

$$R_n = \left\{ \begin{pmatrix} a & a_{12} & a_{13} & \cdots & a_{1n} \\ 0 & a & a_{23} & \cdots & a_{2n} \\ 0 & 0 & a & \cdots & a_{3n} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \cdots & a \end{pmatrix} \middle| a, a_{ij} \in S, n \ge 3 \right\}$$

is not reversible by [6, Example 1.5]. But  $R_n$  is weakly reversible by Proposition 2.3 since any subring of weakly reversible rings is weakly reversible. It is obvious that  $R_4$  is not semicommutative and it can be proved similarly that  $R_n$  is not semicommutative for  $n \geq 5$ .

The next example demonstrates that the condition *n*-by-*n* upper triangular matrix ring  $T_n(R)$  in Proposition 2.3 cannot be weakened to *n*-by-*n* full matrix ring  $M_n(R)$ , where *n* is any integer greater than 1.

**Example 2.7.** If R is a weakly reversible ring and n is any integer greater than 1, then  $M_n(R)$  is not weakly reversible.

For n=2, observe that

but

$$\begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix} = 0.$$
$$\begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix} \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix} \bar{\in} \operatorname{nil}(M_2(R))$$

So  $M_2(R)$  is not weakly reversible. One can augment these matrices in a similar way if n > 2.

Acknowledgement. The authors would like to express their sincere gratitude to the referee for valuable suggestions and many careful comments which led to substantial improvements of this paper.

### ZHAO LIANG AND YANG GANG

#### References

- 1. Cohn P. M., *Reversible rings*, Bull. London Math. Soc. 31 (1999), 641–648.
- Anderson D. D. and Camillo V., Semigroups and rings whose zero products commute, Comm. Algebra 27(6) (1999), 2847–2852.
- Krempa J. and Niewieczerzal D., Rings in which annihilators are ideals and their application to semigroup rings, Bull. Acad. Polon. Sci. Ser. Sci., Math. Astronom, Phys. 25 (1977), 851–856.
- Shin G., Prime ideals and sheaf representation of a pseudo symmetric ring, Trans. Amer. Math. Soc. 184 (1973), 43–60.
- Lambek J., On the representation of modules by sheaves of factor modules, Canad. Math. Bull. 14(3) (1971), 359–368.
- Kim N. K. and Lee Y., Extensions of reversible rings, J. Pure Appl. Algebra 185 (2003), 207–223.
- 7. Marks G., Reversible and symmetric rings, J. Pure Appl. Algebra 174 (2002), 311–318.
- Anderson D. D. and Camillo V., Armendariz rings and Gaussian rings, Comm. Algebra 26(7) (1998), 2265–2272.
- 9. Marks G., A taxonomy of 2-primal rings, J. Algebra 266(2) (2003), 494-520.

Zhao Liang, School of Science, Jiangxi University of Science and Technology, Jiangxi 341000, China, *e-mail*: zhaoliang.nwnu@126.com

Yang Gang, School of Mathematics, Physics Engineering, Lanzhou Jiaotong University, Gansu 730070, China