

ON PSEUDO-SEQUENCE-COVERING π -IMAGES OF LOCALLY SEPARABLE METRIC SPACES

NGUYEN VAN DUNG

ABSTRACT. In this paper, we characterize pseudo-sequence-covering π -images of locally separable metric spaces by means of fcs -covers and point-star networks. We also investigate pseudo-sequence-covering π - s -images of locally separable metric spaces.

1. INTRODUCTION

Determining what spaces the images of “nice” spaces under “nice” mappings are is one of the central questions of general topology [3]. In the past, some noteworthy results on images of metric spaces have been obtained [9, 15]. Recently, π -images of metric spaces have attracted attention again [4, 5, 7, 11, 16]. It is known that a space is a pseudo-sequence-covering π -image of a metric space (resp. separable metric space) if and only if it has a point-star network of fcs -covers (resp. countable fcs -covers) [4, 5]. This leads us to investigate pseudo-sequence-covering π -images of locally separable metric spaces. That is, we have the following question.

Question 1.1. *How are pseudo-sequence-covering π -images of locally separable metric spaces characterized?*

On the other hand, pseudo-sequence-covering π - s -images of metric spaces have been characterized by means of point-star networks of point-countable fcs -covers (see [11], for example). This leads us to consider the following question.

Question 1.2. *How are pseudo-sequence-covering π - s -images of locally separable metric spaces characterized?*

Taking these questions into account, we characterize pseudo-sequence-covering π -images of locally separable metric spaces by means of fcs -covers and point-star networks. Then we give a complete answer to Question 1.1. As the application

Received May 20, 2007.

2000 *Mathematics Subject Classification.* Primary 54E40; Secondary 54C10, 54E99, 54D55, 54D65.

Key words and phrases. π -map; s -map; network; k -cover; cfp -cover; fcs -cover; cs^* -cover; pseudo-sequence-covering; subsequence-covering; sequentially-quotient.

Supported in part by the National Natural Science Foundation of Viet Nam.

of this result, we get a characterization of pseudo-sequence-covering π - s -images of locally separable metric spaces to answer Question 1.2.

Throughout this paper, all spaces are assumed to be Hausdorff, all mappings are assumed continuous and onto, a convergent sequence includes its limit point, \mathbb{N} denotes the set of all natural numbers. Let $f : X \rightarrow Y$ be a mapping, $x \in X$, and let \mathcal{P} be a collection of subsets of X , we denote $\text{st}(x, \mathcal{P}) = \bigcup\{P \in \mathcal{P} : x \in P\}$, $\bigcup \mathcal{P} = \bigcup\{P : P \in \mathcal{P}\}$, $(\mathcal{P})_x = \{P \in \mathcal{P} : x \in P\}$ and $f(\mathcal{P}) = \{f(P) : P \in \mathcal{P}\}$. We say that a convergent sequence $\{x_n : n \in \mathbb{N}\}$ converging to x is *eventually* (resp. *frequently*) in A if $\{x_n : n \geq n_0\} \cup \{x\} \subset A$ for some $n_0 \in \mathbb{N}$ (resp. $\{x_{n_k} : k \in \mathbb{N}\} \cup \{x\} \subset A$ for some subsequence $\{x_{n_k}\}$ of $\{x_n\}$). Note that some notions are different in different references, and some different notions in different references are coincident. Please, terms which are not defined here, see [2, 15].

2. MAIN RESULTS

Let \mathcal{P} be a collection of subsets of a space X and let K be a subset of X .

\mathcal{P} is *point-countable* [15] if every point of X meets only countably many members of \mathcal{P} .

For each $x \in X$, \mathcal{P} is a *network at x* [8] if $x \in P$ for every $P \in \mathcal{P}$, and if $x \in U$ with U open in X , there exists $P \in \mathcal{P}$ such that $x \in P \subset U$.

\mathcal{P} is a *k -cover for K in X* , if for each compact subset H of K , there exists a finite subfamily \mathcal{F} of \mathcal{P} such that $H \subset \bigcup \mathcal{F}$. When $K = X$, a *k -cover for K in X* is a *k -cover for X* .

\mathcal{P} is a *cfp-cover for K in X* if for each compact subset H of K , there exists a finite subfamily \mathcal{F} of \mathcal{P} such that $H \subset \bigcup\{C_F : F \in \mathcal{F}\}$ where C_F is closed and $C_F \subset F$ for every $F \in \mathcal{F}$. Note that such \mathcal{F} is a *full cover* in the sense of [1], and if K is closed, \mathcal{F} is a *cfp-cover for K in the sense of [8]*. When $K = X$, a *cfp-cover for K in X* is a *cfp-cover for X* [16].

\mathcal{P} is an *fcs-cover for K in X* if for each convergent sequence S converging to x in K , there exists a finite subfamily \mathcal{F} of $(\mathcal{P})_x$ such that S is eventually in $\bigcup \mathcal{F}$. When $K = X$, an *fcs-cover for K in X* is an *fcs-cover of X* [4], or an *sfp-cover for X* [11], or a *wcs-cover* [5].

\mathcal{P} is a *cs*-cover for K in X* , if for each convergent sequence S in K , S is frequently in some $P \in \mathcal{P}$. When $K = X$, a *cs*-cover for K in X* is a *cs*-cover for X* [16].

A *k -cover* (resp. *cfp-cover*, *fcs-cover*, *cs*-cover*) for K in X is also called a *k -cover* (resp. *cfp-cover*, *fcs-cover*, *cs*-cover*) *in X for K* , and a *k -cover* (resp. *cfp-cover*, *fcs-cover*, *cs*-cover*) for X is abbreviated to a *k -cover* (resp. *cfp-cover*, *fcs-cover*, *cs*-cover*).

It is clear that if \mathcal{P} is a *k -cover* (resp. *cfp-cover*, *fcs-cover*, *cs*-cover*), then \mathcal{P} is a *k -cover* (resp. *cfp-cover*, *fcs-cover*, *cs*-cover*) for K in X .

Remark. The following statements hold.

1. closed *k -cover for K in X* \implies *cfp-cover for K in X* \implies *k -cover for K in X* ,

2. *cfp*-cover for K in $X \implies fcs$ -cover for K in $X \implies cs^*$ -cover for K in X .

For each $n \in \mathbb{N}$, let \mathcal{P}_n be a cover for X . $\{\mathcal{P}_n : n \in \mathbb{N}\}$ is a *refinement sequence* for X , if \mathcal{P}_{n+1} is a refinement of \mathcal{P}_n for each $n \in \mathbb{N}$. A refinement sequence for X is a *refinement* of X in the sense of [3].

Let $\{\mathcal{P}_n : n \in \mathbb{N}\}$ is be refinement sequence for X . $\{\mathcal{P}_n : n \in \mathbb{N}\}$ is a *point-star network* for X , if $\{st(x, \mathcal{P}_n) : n \in \mathbb{N}\}$ is a network at x for each $x \in X$. A point-star network for X is a σ -*strong network* for X in the sense of [16], and, without the assumption of a refinement sequence, a *point-star network* in the sense of [12]. It is easy to see that if each \mathcal{P}_n is countable, every members of \mathcal{P}_n can be chosen closed in X .

Let $\{\mathcal{P}_n : n \in \mathbb{N}\}$ be a point-star network for a space X . For every $n \in \mathbb{N}$, put $\mathcal{P}_n = \{P_\alpha : \alpha \in A_n\}$, and A_n is endowed with discrete topology. Put

$$M = \{a = (\alpha_n) \in \prod_{n \in \mathbb{N}} A_n : \{P_{\alpha_n} : n \in \mathbb{N}\}$$

forms a network at some point x_a in $X\}$.

Then M , which is a subspace of the product space $\prod_{n \in \mathbb{N}} A_n$, is a metric space with a metric d described as follows.

Let $a = (\alpha_n), b = (\beta_n) \in M$. If $a = b$, then $d(a, b) = 0$. If $a \neq b$, then $d(a, b) = 1/(\min\{n \in \mathbb{N} : \alpha_n \neq \beta_n\})$.

Define $f : M \rightarrow X$ by choosing $f(a) = x_a$, then f is a mapping, and $(f, M, X, \{\mathcal{P}_n\})$ is a *Ponomarev's system* [16], and without the assumption of a refinement sequence in the notion of point-star networks, $(f, M, X, \{\mathcal{P}_n\})$ is a *Ponomarev's system* in the sense of [12].

Let $f : X \rightarrow Y$ be a mapping; Then,

f is a π -*mapping* [4] if for every $y \in Y$ and for every neighborhood U of y in Y , $d(f^{-1}(y), X - f^{-1}(U)) > 0$, where X is a metric space with a metric d .

f is an s -*mapping* [11], if for each $y \in Y$, $f^{-1}(y)$ is a separable subset of X .

f is a π - s -*mapping* [11], if f is both π -mapping and s -mapping.

f is a *pseudo-sequence-covering mapping* [3], if every convergent sequence of Y is the image of some compact subset of X .

f is a *subsequence-covering mapping* [3], if for every convergent sequence S of Y , there is a compact subset K of X such that $f(K)$ is a subsequence of S .

f is a *sequentially-quotient mapping* [3], if for every convergent sequence S of Y , there is a convergent sequence L of X such that $f(L)$ is a subsequence of S .

f is a *quotient mapping* [14], if U is open in Y whenever $f^{-1}(U)$ is open in X .

f is a *pseudo-open mapping* [9], if $y \in \text{int} f(U)$ whenever $f^{-1}(y) \subset U$ with U open in X . A pseudo-open mapping is a *hereditarily quotient mapping* in the sense of [2].

Let X be a space and let A be a subset of X . A is *sequential open* [16], if for each $x \in A$ and each convergent sequence S converging to x , S is eventually in A . X is a *sequential space* [16], if every sequential open subset of X is open in X . X is a *Fréchet space*, if for each $x \in \bar{A}$, there exists a sequence in A converging to x .

For a mapping $f : X \rightarrow Y$, f is a pseudo-sequence-covering or sequentially-quotient \implies a f is subsequence-covering. Also, a f is quotient if and only if a f is subsequence-covering such that Y is sequential [17].

Lemma 2.1. *Let \mathcal{P} be a countable cover for a convergent sequence S in a space X . Then the following propositions are equivalent.*

1. \mathcal{P} is a *cfp*-cover for S in X ,
2. \mathcal{P} is an *fcs*-cover for S in X ,
3. \mathcal{P} is a *cs**-cover for S in X .

Proof. (1) \implies (2) \implies (3). Obviously.

(3) \implies (1). Let H be a compact subset of S . We can assume that H is a subsequence of S . Since \mathcal{P} is countable, put $(\mathcal{P})_x = \{P_n : n \in \mathbb{N}\}$ where x is the limit point of S . Then H is eventually in $\bigcup_{n \leq k} P_n$ for some $k \in \mathbb{N}$. If not, then for any $k \in \mathbb{N}$, H is not eventually in $\bigcup_{n \leq k} P_n$. So, for every $k \in \mathbb{N}$, there exists $x_{n_k} \in S - \bigcup_{n \leq k} P_n$. We may assume $n_1 < n_2 < \dots < n_{k-1} < n_k < n_{k+1} < \dots$. Put $H' = \{x_{n_k} : k \in \mathbb{N}\} \cup \{x\}$, then H' is a subsequence of S . Since \mathcal{P} is a *cs**-cover for S in X , there exists $m \in \mathbb{N}$ such that H' is frequently in P_m . This contradicts the construction of H' . So H is eventually in $\bigcup_{n \leq k} P_n$ for some $k \in \mathbb{N}$. It implies that \mathcal{P} is a *cfp*-cover for S in X . \square

Lemma 2.2. *Let $f : X \rightarrow Y$ be a mapping.*

1. If \mathcal{P} is a *k*-cover in X for a compact set K , then $f(\mathcal{P})$ is a *k*-cover for $f(K)$ in Y .
2. If \mathcal{P} is a *cfp*-cover in X for a compact set K , then $f(\mathcal{P})$ is a *cfp*-cover for $f(K)$ in Y .

Proof. (1). Let H be a compact subset of $f(K)$. Then $G = f^{-1}(H) \cap K$ is a compact subset of K and $f(G) = H$. Since \mathcal{P} is a *k*-cover for K in X , there is a finite subfamily \mathcal{F} of \mathcal{P} such that $G \subset \bigcup \mathcal{F}$. Hence $f(\mathcal{F})$ is a finite subfamily of $f(\mathcal{P})$ such that $H \subset \bigcup f(\mathcal{F})$. It implies that $f(\mathcal{P})$ is a *k*-cover for $f(K)$ in Y .

(2). Let H be a compact subset of $f(K)$. Then $L = f^{-1}(H) \cap K$ is a compact subset of K satisfying $f(L) = H$. Since \mathcal{P} is a *cfp*-cover for K in X , there is a finite subfamily \mathcal{F} of \mathcal{P} such that $L \subset \bigcup \{C_F : F \in \mathcal{F}\}$ where $C_F \subset F$, and C_F is closed for every $F \in \mathcal{F}$. Because L is compact, every C_F can be chosen compact. It implies that every $f(C_F)$ is closed (in fact, every $f(C_F)$ is compact), and $f(C_F) \subset f(F)$. We get that $H = f(L) \subset \bigcup \{f(C_F) : F \in \mathcal{F}\}$, and $f(\mathcal{F})$ is a finite subfamily of \mathcal{P} . Then \mathcal{P} is a *cfp*-cover for $f(K)$ in Y . \square

Theorem 2.3. *The following propositions are equivalent for a space X*

1. X is a pseudo-sequence-covering π -image of a locally separable metric space,
2. X has a cover $\{X_\lambda : \lambda \in \Lambda\}$, where each X_λ has a refinement sequence $\{\mathcal{P}_{\lambda,n} : n \in \mathbb{N}\}$ of countable covers for X_λ satisfying the following conditions:
 - (a) For each $x \in U$ with U open in X , there is $n \in \mathbb{N}$ such that

$$\bigcup \{\text{st}(x, \mathcal{P}_{\lambda,n}) : \lambda \in \Lambda \text{ with } x \in X_\lambda\} \subset U,$$

- (b) For each convergent sequence S of X , there is a finite subset Λ_S of Λ such that S has a finite compact cover $\{S_\lambda : \lambda \in \Lambda_S\}$, and, for each $\lambda \in \Lambda_S$ and $n \in \mathbb{N}$, $\mathcal{P}_{\lambda,n}$ is an fcs-cover for S_λ in X_λ .

Proof. (1) \implies (2). Let $f : M \rightarrow X$ be a pseudo-sequence-covering π -mapping from a locally separable metric space M with a metric d onto X . Since M is a locally separable metric space, $M = \bigoplus_{\lambda \in \Lambda} M_\lambda$ where each M_λ is a separable metric space by [2, 4.4.F]. For each $\lambda \in \Lambda$, let D_λ be a countable dense subset of M_λ , and put $f_\lambda = f|_{M_\lambda}$ and $X_\lambda = f_\lambda(M_\lambda)$. For each $a \in M_\lambda$ and $n \in \mathbb{N}$, put $B(a, 1/n) = \{b \in M_\lambda : d(a, b) < 1/n\}$, $\mathcal{B}_{\lambda,n} = \{B(a, 1/n) : a \in D_\lambda\}$, and $\mathcal{P}_{\lambda,n} = f_\lambda(\mathcal{B}_{\lambda,n})$. It is clear that $\{\mathcal{P}_{\lambda,n} : n \in \mathbb{N}\}$ is a cover sequence of countable covers for X_λ and $\mathcal{P}_{\lambda,n+1}$ is a refinement of $\mathcal{P}_{\lambda,n}$ for every $n \in \mathbb{N}$. We only need to prove that conditions (a) and (b) are satisfied.

Condition (a): For each $x \in U$ with U open in X . Since f is a π -mapping, $d(f^{-1}(x), M - f^{-1}(U)) > 2/(n - 1)$ for some $n \in \mathbb{N}$. Then, for each $\lambda \in \Lambda$ with $x \in X_\lambda$, we get

$$d(f_\lambda^{-1}(x), M_\lambda - f_\lambda^{-1}(U_\lambda)) > 2/(n - 1)$$

where $U_\lambda = U \cap X_\lambda$. Let $a \in D_\lambda$ and $x \in f_\lambda(B(a, 1/n)) \in \mathcal{P}_{\lambda,n}$. We shall prove that $B(a, 1/n) \subset f_\lambda^{-1}(U_\lambda)$. In fact, if $B(a, 1/n) \not\subset f_\lambda^{-1}(U_\lambda)$, then pick $b \in B(a, 1/n) - f_\lambda^{-1}(U_\lambda)$. Note that $f_\lambda^{-1}(x) \cap B(a, 1/n) \neq \emptyset$, pick $c \in f_\lambda^{-1}(x) \cap B(a, 1/n)$, then

$$d(f_\lambda^{-1}(x), M_\lambda - f_\lambda^{-1}(U_\lambda)) \leq d(c, b) \leq d(c, a) + d(a, b) < 2/n < 2/(n - 1).$$

It is a contradiction. So $B(a, 1/n) \subset f_\lambda^{-1}(U_\lambda)$, thus $f_\lambda(B(a, 1/n)) \subset U_\lambda$. Then $\text{st}(x, \mathcal{P}_{\lambda,n}) \subset U_\lambda$. It implies that

$$\bigcup \{\text{st}(x, \mathcal{P}_{\lambda,n}) : \lambda \in \Lambda \text{ with } x \in X_\lambda\} \subset U.$$

Condition (b): For each convergent sequence S of X , since a f is pseudo-sequence-covering, $S = f(K)$ for some compact subset K of M . By compactness of K , $K_\lambda = K \cap M_\lambda$ is compact and $\Lambda_S = \{\lambda \in \Lambda : K_\lambda \neq \emptyset\}$ is finite. For each $\lambda \in \Lambda_S$, put $S_\lambda = f(K_\lambda)$, then $\{S_\lambda : \lambda \in \Lambda_S\}$ is a finite compact cover for S . For each $n \in \mathbb{N}$, since $\mathcal{B}_{\lambda,n}$ is a cfp-cover for K_λ in M_λ , $\mathcal{P}_{\lambda,n}$ is a cfp-cover for S_λ in X_λ by Lemma 2.2. It follows from Lemma 2.1 that $\mathcal{P}_{\lambda,n}$ is an fcs-cover for S_λ in X_λ .

(2) \implies (1). For each $\lambda \in \Lambda$, let $x \in U_\lambda$ with U_λ open in X_λ . We get that $U_\lambda = U \cap X_\lambda$ with some U open in X . Since $\bigcup \{\text{st}(x, \mathcal{P}_{\lambda,n}) : \lambda \in \Lambda \text{ with } x \in X_\lambda\} \subset U$ for some $n \in \mathbb{N}$, $\text{st}(x, \mathcal{P}_{\lambda,n}) \subset U_\lambda$. It implies $\{\mathcal{P}_{\lambda,n} : n \in \mathbb{N}\}$ is a point-star network for X_λ . Then the Ponomarev's system $(f_\lambda, M_\lambda, X_\lambda, \{\mathcal{P}_{\lambda,n}\})$ exists. Since each $\mathcal{P}_{\lambda,n}$ is countable, M_λ is a separable metric space with a metric d_λ described as follows.

Let $a = (\alpha_n), b = (\beta_n) \in M_\lambda$. If $a = b$, then $d_\lambda(a, b) = 0$. If $a \neq b$, then $d_\lambda(a, b) = 1/(\min\{n \in \mathbb{N} : \alpha_n \neq \beta_n\})$.

Put $M = \bigoplus_{\lambda \in \Lambda} M_\lambda$ and define $f : M \rightarrow X$ by choosing $f(a) = f_\lambda(a)$ for every $a \in M_\lambda$ with some $\lambda \in \Lambda$. Then f is a mapping and M is a locally separable metric space with a metric d as follows.

Let $a, b \in M$. If $a, b \in M_\lambda$ for some $\lambda \in \Lambda$, then $d(a, b) = d_\lambda(a, b)$. Otherwise, $d(a, b) = 1$. We only need to prove that f is a pseudo-sequence-covering π -mapping.

(a) f is a π -mapping. Let $x \in U$ with U open in X , then

$$\bigcup \{st(x, \mathcal{P}_{\lambda,n}) : \lambda \in \Lambda \text{ with } x \in X_\lambda\} \subset U$$

for some $n \in \mathbb{N}$. So, for each $\lambda \in \Lambda$ with $x \in X_\lambda$, we get

$$st(x, \mathcal{P}_{\lambda,n}) \subset U_\lambda$$

where $U_\lambda = U \cap X_\lambda$. It implies that

$$d_\lambda(f_\lambda^{-1}(x), M_\lambda - f_\lambda^{-1}(U_\lambda)) \geq 1/n.$$

In fact, if $a = (\alpha_k) \in M_\lambda$ such that $d_\lambda(f_\lambda^{-1}(x), a) < 1/n$, then there is $b = (\beta_k) \in f_\lambda^{-1}(x)$ such that $d_\lambda(a, b) < 1/n$. So $\alpha_k = \beta_k$ if $k \leq n$. Note that $x \in P_{\beta_n} \subset st(x, \mathcal{P}_{\lambda,n}) \subset U_\lambda$. Then

$$f_\lambda(a) \in P_{\alpha_n} = P_{\beta_n} \subset st(x, \mathcal{P}_{\lambda,n}) \subset U_\lambda.$$

Hence $a \in f_\lambda^{-1}(U_\lambda)$. It implies that $d_\lambda(f_\lambda^{-1}(x), a) \geq 1/n$ if $a \in M_\lambda - f_\lambda^{-1}(U_\lambda)$. So

$$d_\lambda(f_\lambda^{-1}(x), M_\lambda - f_\lambda^{-1}(U_\lambda)) \geq 1/n.$$

Therefore

$$\begin{aligned} d(f^{-1}(x), M - f^{-1}(U)) &= \inf\{d(a, b) : a \in f^{-1}(x), b \in M - f^{-1}(U)\} \\ &= \min\{1, \inf\{d_\lambda(a, b) : a \in f_\lambda^{-1}(x), b \in M_\lambda - f^{-1}(U_\lambda), \lambda \in \Lambda\}\} \geq 1/n > 0. \end{aligned}$$

It implies that f is a π -mapping.

(b) f is pseudo-sequence-covering. For each convergent sequence S of X , there is a finite subset Λ_S of Λ such that S has a finite compact cover $\{S_\lambda : \lambda \in \Lambda_S\}$ and for each $\lambda \in \Lambda_S$ and $n \in \mathbb{N}$, $\mathcal{P}_{\lambda,n}$ is an fcs -cover for S_λ in X_λ . By Lemma 2.1 $\mathcal{P}_{\lambda,n}$ is a cfp -cover for S_λ in X_λ . It follows from Lemma 13 in [12] that $S_\lambda = f_\lambda(K_\lambda)$ with some compact subset K_λ of M_λ . Put $K = \bigcup\{K_\lambda : \lambda \in \Lambda_S\}$, then K is a compact subset of M and $f(K) = S$. It implies that f is a pseudo-sequence-covering. \square

- Remark.**
1. For each $\lambda \in \Lambda$, $\{\mathcal{P}_{\lambda,n} : n \in \mathbb{N}\}$ is a point-star network for X_λ .
 2. Since each $\mathcal{P}_{\lambda,n}$ is countable, every member of $\mathcal{P}_{\lambda,n}$ can be chosen closed in X_λ . Hence, it is possible to replace the prefix “ fcs –” in (b) of Theorem 2.3.(2) by “ k –”, “ cfp –”, or “ cs^* –”

By [2, 2.4.F, 2.4.G], [3, Proposition 2.1], and Theorem 2.3, we get a characterization of pseudo-sequence-covering quotient (resp. pseudo-open) π -images of locally separable metric spaces as follows.

Corollary 2.4. *The following propositions are equivalent:*

1. *a space X is a pseudo-sequence-covering quotient (resp. pseudo-open) π -image of a locally separable metric space,*

2. a space X is a sequential (resp. Fréchet) space having a cover $\{X_\lambda : \lambda \in \Lambda\}$, where each X_λ has a refinement sequence $\{\mathcal{P}_{\lambda,n} : n \in \mathbb{N}\}$ of countable covers for X_λ satisfying conditions (a) and (b) in Theorem 2.3.(2).

In the next, we investigate pseudo-sequence-covering π - s -images of locally separable metric spaces.

Corollary 2.5. *The following propositions are equivalent:*

1. a space X is a pseudo-sequence-covering π - s -image of a locally separable metric space,
2. a space X has a point-countable cover $\{X_\lambda : \lambda \in \Lambda\}$, where each X_λ has a refinement sequence $\{\mathcal{P}_{\lambda,n} : n \in \mathbb{N}\}$ of countable covers for X_λ satisfying conditions (a) and (b) in Theorem 2.3.(2).

Proof. (1) \implies (2). By using notations and arguments in proof (1) \implies (2) of Theorem 2.3 again, X has a cover $\{X_\lambda : \lambda \in \Lambda\}$, where each X_λ has a refinement sequence $\{\mathcal{P}_{\lambda,n} : n \in \mathbb{N}\}$ of countable covers for X_λ satisfying conditions (a) and (b) in Theorem 2.3.(2). It suffices to prove that $\{X_\lambda : \lambda \in \Lambda\}$ is point-countable. For each $x \in X$, since f is an s -mapping, $f^{-1}(x)$ is separable in M . Then $f^{-1}(x)$ meets only countably many M_λ 's. It implies that x meets only countably many X_λ 's, i.e., $\{X_\lambda : \lambda \in \Lambda\}$ is point-countable.

(2) \implies (1). By using notations and arguments in proof (2) \implies (1) of Theorem 2.3 again, X is a pseudo-sequence-covering π -image of a locally separable metric space under the mapping f . We shall prove that f is also an s -mapping. For each $x \in X$, since $\{X_\lambda : \lambda \in \Lambda\}$ is point-countable, $\Lambda_x = \{\lambda \in \Lambda : x \in X_\lambda\}$ is countable. Note that each M_λ is separable metric, $f_\lambda^{-1}(x)$ is separable. It implies that $f^{-1}(x) = \bigcup \{f_\lambda^{-1}(x) : \lambda \in \Lambda_x\}$ is separable, i.e., f is an s -mapping. \square

Similar to Corollary 2.4, we get the following.

Corollary 2.6. *The following propositions are equivalent:*

1. a space X is a pseudo-sequence-covering quotient (resp. pseudo-open) π - s -image of a locally separable metric space,
2. a space X is a sequential (resp. Fréchet) space having a point-countable cover $\{X_\lambda : \lambda \in \Lambda\}$, where each X_λ has a refinement sequence $\{\mathcal{P}_{\lambda,n} : n \in \mathbb{N}\}$ of countable covers for X_λ satisfying conditions (a) and (b) in Theorem 2.3.(2).

Acknowledgment. We would like to thank Professor Y. Ge for helpful comment on this article.

REFERENCES

1. Chen H., *Compact-covering maps and k -networks*, Proc. Amer. Math. Soc., **131** (2002), 2623–2632.
2. Engelking R., *General topology*, PWN-Polish Scientific Publishers, Warszawa 1977.
3. Ge Y., *On compact images of locally separable metric spaces*, Topology Proc., **27** (1) (2003), 351–360.
4. ———, *On π -images of metric spaces*, APN Acta Math., **22** (2006), 209–215.

5. ———, *On pseudo-sequence-covering π -images of metric spaces*, *Matematicki Vesnik*, **57** (2005), 113–120.
6. ———, *Spaces with countable sn -networks*, *Comment. Math. Univ. Carolinae*, **45** (1) (2004), 169–176.
7. Ge Y., and Gu J. S., *On π -images of separable metricspaces*, *Math. Sci.*, **10** (2004), 65–71.
8. Ge Y., and Shen J., *Networks in Ponomarev-systems*, *Math. Sci.*, **11** (2005), 25–29.
9. Gruenhagen G., Michael E. and Tanaka Y., *Spaces determined by point-countable covers*, *Pacific J. Math.*, **113** (2) (1984), 303–332.
10. Ikeda Y., Liu C. and Tanaka Y., *Quotient compact images of metric spaces, and related matters*, *Topology Appl.*, **122** (2002), 237–252.
11. Li Z., *On π - s -images of metric spaces*, *Int. J. Math. Sci.*, **7** (2005), 1101–1107.
12. Lin S. and Yan P., *Notes on cfp -covers*, *Comment. Math. Univ. Carolinae*, **44** (2) (2003), 295–306.
13. ———, *Sequence-covering maps of metric spaces*, *Topology Appl.*, **109** (2001), 301–314.
14. Michael E., *N_0 -spaces*, *J. Math. Mech.*, **15** (1966), 983–1002.
15. Tanaka Y., *Theory of k -network II*, *Q & A in General Top.*, **19** (2001), 27–46.
16. Tanaka Y. and Y. Ge, *Around quotient compact images of metric spaces, and symmetric spaces*, *Houston J. Math.*, **32** (1) (2006), 99–117.
17. Tanaka Y. and Z. Li, *Certain covering-maps and k -networks, and related matters*, *Topology Proc.*, **27** (2003), 317–334.

Nguyen Van Dung, Department of Mathematics, Pedagogical University of Dong Thap, Viet Nam, *current address*: 85 Nguyen Phong Sac, Hung Dung street, Vinh city, Nghe An province, *e-mail*: nguyendungtc@yahoo.com