

## SIGNED STAR $(j, k)$ -DOMATIC NUMBER OF A GRAPH

S. M. SHEIKHOESLAMI AND L. VOLKMANN

ABSTRACT. Let  $G$  be a simple graph without isolated vertices with edge set  $E(G)$ , and let  $j$  and  $k$  be two positive integers. A function  $f: E(G) \rightarrow \{-1, 1\}$  is said to be a signed star  $j$ -dominating function on  $G$  if  $\sum_{e \in E(v)} f(e) \geq j$  for every vertex  $v$  of  $G$ , where  $E(v) = \{uv \in E(G) \mid u \in N(v)\}$ . A set  $\{f_1, f_2, \dots, f_d\}$  of distinct signed star  $j$ -dominating functions on  $G$  with the property that  $\sum_{i=1}^d f_i(e) \leq k$  for each  $e \in E(G)$ , is called a signed star  $(j, k)$ -dominating family (of functions) on  $G$ . The maximum number of functions in a signed star  $(j, k)$ -dominating family on  $G$  is the signed star  $(j, k)$ -domatic number of  $G$  denoted by  $d_{SS}^{(j,k)}(G)$ .

In this paper we study properties of the signed star  $(j, k)$ -domatic number of a graph  $G$ . In particular, we determine bounds on  $d_{SS}^{(j,k)}(G)$ . Some of our results extend those ones given by Atapour, Sheikholeslami, Ghameslou and Volkmann [1] for the signed star domatic number, Sheikholeslami and Volkmann [5] for the signed star  $(k, k)$ -domatic number and Sheikholeslami and Volkmann [4] for the signed star  $k$ -domatic number.

### 1. INTRODUCTION

Let  $G$  be a graph with vertex set  $V(G)$  and edge set  $E(G)$ . We use [2] for terminology and notation which are not defined here and consider simple graphs without isolated vertices only. The integers  $n = |V(G)|$  and  $m = |E(G)|$  are the *order* and the *size* of the graph  $G$ , respectively. For every vertex  $v \in V(G)$ , the *open neighborhood*  $N(v)$  of  $v$  is the set  $\{u \in V(G) \mid uv \in E(G)\}$ , and the *closed neighborhood* of  $v$  is the set  $N[v] = N(v) \cup \{v\}$ . The *degree* of a vertex  $v$  is  $d(v) = |N(v)|$ . The *minimum* and *maximum degree* of a graph  $G$  are denoted by  $\delta(G)$  and  $\Delta(G)$ , respectively. The *complement*  $\overline{G}$  of a graph  $G$  is the graph with vertex set  $V(G)$  such that two vertices are adjacent in  $\overline{G}$  if and only if these vertices are not adjacent in  $G$ .

The *open neighborhood*  $N_G(e)$  of an edge  $e \in E(G)$  is the set of all edges adjacent to  $e$ . Its *closed neighborhood* is  $N_G[e] = N_G(e) \cup \{e\}$ . For a function  $f: E(G) \rightarrow \{-1, 1\}$  and a subset  $S$  of  $E(G)$ , we define  $f(S) = \sum_{e \in S} f(e)$ . The *edge-neighborhood*  $E_G(v) = E(v)$  of a vertex  $v \in V(G)$  is the set of all edges

---

Received September 4, 2012.

2010 *Mathematics Subject Classification*. Primary 05C69.

*Key words and phrases*. Signed star  $(j, k)$ -domatic number; Signed star domatic number; Signed star  $j$ -dominating function; Signed star dominating function; Signed star  $j$ -domination number; Signed star domination number.

incident with the vertex  $v$ . For each vertex  $v \in V(G)$ , we also define  $f(v) = \sum_{e \in E_G(v)} f(e)$ .

Let  $j$  be a positive integer. A function  $f: E(G) \rightarrow \{-1, 1\}$  is called a *signed star  $j$ -dominating function* (SSjDF) on  $G$  if  $f(v) \geq j$  for every vertex  $v$  of  $G$ . The *signed star  $j$ -domination number* of a graph  $G$  is  $\gamma_{jSS}(G) = \min\{\sum_{e \in E(G)} f(e) \mid f \text{ is a SSjDF on } G\}$ . The signed star  $j$ -dominating function  $f$  on  $G$  with  $f(E(G)) = \gamma_{jSS}(G)$  is called a  $\gamma_{jSS}(G)$ -*function*. As the assumption  $\delta(G) \geq j$  is clearly necessary, we will always assume that satisfy  $\delta(G) \geq j$  while discussing  $\gamma_{jSS}(G)$  all graphs involved. The signed star  $j$ -domination number was introduced by Xu and Li [10] in 2009 and has been studied by several authors (see for instance, [3, 4, 7]). The signed star 1-domination number is the usual signed star domination number, introduced in 2005 by Xu [8]. The signed star domination number was investigated for example, by [3, 6, 9].

Let  $k$  be a further positive integer. A set  $\{f_1, f_2, \dots, f_d\}$  of distinct signed star  $j$ -dominating functions on  $G$  with  $\sum_{i=1}^d f_i(e) \leq k$  for each  $e \in E(G)$ , is called a *signed star  $(j, k)$ -dominating family* (SS(j,k)D family) (of functions) on  $G$ . The maximum number of functions in a signed star  $(j, k)$ -dominating family on  $G$  is the *signed star  $(j, k)$ -domatic number* of  $G$  denoted by  $d_{SS}^{(j,k)}(G)$ . The signed star  $(j, k)$ -domatic number is well-defined and

$$(1) \quad d_{SS}^{(j,k)}(G) \geq 1$$

for all graphs  $G$  with  $\delta(G) \geq j$ , since the set consisting of any signed star  $j$ -dominating function forms a SS(j,k)D family on  $G$ . A  $d_{SS}^{(j,k)}$ -*family* of a graph  $G$  is a SS(j,k)D family containing exactly  $d_{SS}^{(j,k)}(G)$  signed star  $j$ -dominating functions. The signed star (1,1)-domatic number  $d_{SS}^{(1,1)}(G)$  is the usual signed star domatic number  $d_{SS}(G)$  which was introduced by Atapour, Sheikholeslami, Ghameslou and Volkmann [1] in 2010.

Our purpose in this paper is to initiate the study of the signed star  $(j, k)$ -domatic number in graphs. We study basic properties and bounds for the signed star  $(j, k)$ -domatic number  $d_{SS}^{(j,k)}(G)$  of a graph  $G$ . In addition, we derive Nordhaus-Gaddum type results and bounds of the product and the sum of  $\gamma_{jSS}(G)$  and  $d_{SS}^{(j,k)}(G)$ . Many of our results extend those given by Atapour, Sheikholeslami, Ghameslou and Volkmann [1] for the signed star domatic number, Sheikholeslami and Volkmann [5] for the signed star  $(k, k)$ -domatic number and Sheikholeslami and Volkmann [4] for the signed star  $k$ -domatic number.

**Observation 1** ([4]). *Let  $G$  be a graph of size  $m$  with  $\delta(G) \geq j$ . Then  $\gamma_{jSS}(G) = m$  if and only if each edge  $e \in E(G)$  has an endpoint  $u$  such that  $d(u) = j$  or  $d(u) = j + 1$ .*

2. PROPERTIES OF THE SIGNED STAR  $(j, k)$ -DOMATIC NUMBER

**Theorem 2.** *Let  $j, k \geq 1$  be two integers. If  $G$  is a graph of minimum degree  $\delta(G) \geq j$ , then*

$$d_{SS}^{(j,k)}(G) \leq \frac{k\delta(G)}{j}.$$

Moreover, if  $d_{SS}^{(j,k)}(G) = k\delta(G)/j$ , then for each function of any signed star  $(j, k)$ -dominating family  $\{f_1, f_2, \dots, f_d\}$  with  $d = d_{SS}^{(j,k)}(G)$  and for all vertices  $v$  of degree  $\delta(G)$ ,  $\sum_{e \in E_G(v)} f_i(e) = j$  and  $\sum_{i=1}^d f_i(e) = k$  for every  $e \in E_G(v)$ .

*Proof.* Let  $\{f_1, f_2, \dots, f_d\}$  be a signed star  $(j, k)$ -dominating family on  $G$  such that  $d = d_{SS}^{(j,k)}(G)$ . If  $v \in V(G)$  is a vertex of minimum degree  $\delta(G)$ , then it follows that

$$\begin{aligned} d \cdot j &= \sum_{i=1}^d j \leq \sum_{i=1}^d \sum_{e \in E_G(v)} f_i(e) \\ &= \sum_{e \in E_G(v)} \sum_{i=1}^d f_i(e) \\ &\leq \sum_{e \in E_G(v)} k = k \cdot \delta(G), \end{aligned}$$

and this implies the desired upper bound on the signed star  $(j, k)$ -domatic number.

If  $d_{SS}^{(j,k)}(G) = k\delta(G)/j$ , then the two inequalities occurring in the proof become equalities, which leads to the two properties given in the statement.  $\square$

The special cases  $j = k = 1$ ,  $j = 1$  and  $j = k$  in Theorem 2 can be found in [1], [4] and [5], respectively. As an application of Theorem 2, we will prove the following Nordhaus-Gaddum type result.

**Corollary 3.** *Let  $j, k \geq 1$  be integers. If  $G$  is a graph of order  $n$  such that  $\delta(G) \geq j$  and  $\delta(\overline{G}) \geq j$ , then*

$$d_{SS}^{(j,k)}(G) + d_{SS}^{(j,k)}(\overline{G}) \leq \frac{k}{j}(n-1).$$

If  $d_{SS}^{(j,k)}(G) + d_{SS}^{(j,k)}(\overline{G}) = k(n-1)/j$ , then  $G$  is regular.

*Proof.* Since  $\delta(G) \geq j$  and  $\delta(\overline{G}) \geq j$ , it follows from Theorem 2 that

$$\begin{aligned} d_{SS}^{(j,k)}(G) + d_{SS}^{(j,k)}(\overline{G}) &\leq \frac{k\delta(G)}{j} + \frac{k\delta(\overline{G})}{j} \\ &= \frac{k}{j}(\delta(G) + (n - \Delta(G) - 1)) \leq \frac{k}{j}(n-1), \end{aligned}$$

and this is the desired Nordhaus-Gaddum inequality. If  $G$  is not regular, then  $\Delta(G) - \delta(G) \geq 1$ , and the above inequality chain leads to the better bound  $d_{SS}^{(j,k)}(G) + d_{SS}^{(j,k)}(\overline{G}) \leq \frac{k}{j}(n-2)$ . This completes the proof.  $\square$

**Theorem 4.** *Let  $j, k \geq 1$  be integers. If  $v$  is a vertex of a graph  $G$  such that  $d(v)$  is odd and  $j$  is even or  $d(v)$  is even and  $j$  is odd, then*

$$d_{SS}^{(j,k)}(G) \leq \frac{k}{j+1} \cdot d(v).$$

*Proof.* Let  $\{f_1, f_2, \dots, f_d\}$  be a signed star  $(j, k)$ -dominating family on  $G$  such that  $d = d_{SS}^{(j,k)}(G)$ . Assume first that  $d(v)$  is odd and  $j$  is even. The definition yields to  $\sum_{e \in E_G(v)} f_i(e) \geq j$  for each  $i \in \{1, 2, \dots, d\}$ . On the left-hand side of this inequality a sum of an odd number of odd summands occurs. Therefore it is an odd number, and as  $j$  is even, we obtain  $\sum_{e \in E_G(v)} f_i(e) \geq j+1$  for each  $i \in \{1, 2, \dots, d\}$ . It follows that

$$\begin{aligned} k \cdot d(v) &= \sum_{e \in E_G(v)} k \geq \sum_{e \in E_G(v)} \sum_{i=1}^d f_i(e) \\ &= \sum_{i=1}^d \sum_{e \in E_G(v)} f_i(e) \geq \sum_{i=1}^d (j+1) = d(j+1), \end{aligned}$$

and this leads to the desired bound. Assume next that  $d(v)$  is even and  $j$  is odd. Note that  $\sum_{e \in E_G(v)} f_i(e) \geq j$  for each  $i \in \{1, 2, \dots, d\}$ . On the left-hand side of this inequality a sum of an even number of odd summands occurs. Therefore it is an even number, and as  $j$  is odd, we obtain  $\sum_{e \in E_G(v)} f_i(e) \geq j+1$  for each  $i \in \{1, 2, \dots, d\}$ . Now the desired bound follows as above, and the proof is complete.  $\square$

The next result is an immediate consequence of Theorem 4.

**Corollary 5.** *Let  $j, k \geq 1$  be integers. If  $G$  is a graph such that  $\delta(G)$  is odd and  $j$  is even or  $\delta(G)$  is even and  $j$  is odd, then*

$$d_{SS}^{(j,k)}(G) \leq \frac{k}{j+1} \cdot \delta(G).$$

As an application of Corollary 5, we will improve the Nordhaus-Gaddum bound in Corollary 3 for many cases.

**Theorem 6.** *Let  $j, k \geq 1$  be two integers and let  $G$  be a graph of order  $n$  such that  $\delta(G) \geq j$  and  $\delta(\overline{G}) \geq j$ . If  $\Delta(G) - \delta(G) \geq 1$  or  $j$  is odd or  $j$  is even and  $\delta(G)$  is odd or  $j, \delta(G)$  and  $n$  are even, then*

$$d_{SS}^{(j,k)}(G) + d_{SS}^{(j,k)}(\overline{G}) < \frac{k}{j}(n-1).$$

*Proof.* If  $\Delta(G) - \delta(G) \geq 1$ , then Corollary 3 implies the desired bound. Thus assume now that  $G$  is  $\delta(G)$ -regular.

Case 1. Assume that  $j$  is odd. If  $\delta(G)$  is even, then from Theorem 2 and Corollary 5 it follows that

$$\begin{aligned} d_{SS}^{(j,k)}(G) + d_{SS}^{(j,k)}(\overline{G}) &\leq \frac{k}{j+1}\delta(G) + \frac{k}{j}\delta(\overline{G}) \\ &< \frac{k}{j}(\delta(G) + (n - \delta(G) - 1)) \\ &= \frac{k}{j}(n - 1). \end{aligned}$$

If  $\delta(G)$  is odd, then  $n$  is even and thus  $\delta(\overline{G}) = n - \delta(G) - 1$  is even. Combining Theorem 2 and Corollary 5, we find that

$$\begin{aligned} d_{SS}^{(j,k)}(G) + d_{SS}^{(j,k)}(\overline{G}) &\leq \frac{k}{j}\delta(G) + \frac{k}{j+1}\delta(\overline{G}) \\ &< \frac{k}{j}(\delta(G) + (n - \delta(G) - 1)) \\ &= \frac{k}{j}(n - 1), \end{aligned}$$

and this completes the proof of Case 1.

Case 2. Assume that  $j$  is even. If  $\delta(G)$  is odd, then from Theorem 2 and Corollary 5 it follows that

$$d_{SS}^{(j,k)}(G) + d_{SS}^{(j,k)}(\overline{G}) \leq \frac{k}{j+1}\delta(G) + \frac{k}{j}(n - \delta(G) - 1) < \frac{k}{j}(n - 1).$$

If  $\delta(G)$  is even and  $n$  is even, then  $\delta(\overline{G}) = n - \delta(G) - 1$  is odd, and we obtain the desired bound as above.  $\square$

**Theorem 7.** *Let  $j, k \geq 1$  be integers. If  $G$  is a graph such that  $k$  is odd and  $d_{SS}^{(j,k)}(G)$  is even or  $k$  is even and  $d_{SS}^{(j,k)}(G)$  is odd, then*

$$d_{SS}^{(j,k)}(G) \leq \frac{k-1}{j} \cdot \delta(G).$$

*Proof.* Let  $\{f_1, f_2, \dots, f_d\}$  be a signed star  $(j, k)$ -dominating family on  $G$  such that  $d = d_{SS}^{(j,k)}(G)$ . Assume first that  $k$  is odd and  $d$  is even. If  $e \in E(G)$  is an arbitrary edge, then  $\sum_{i=1}^d f_i(e) \leq k$ . On the left-hand side of this inequality a sum of an even number of odd summands occurs. Therefore, it is an even number, and as  $k$  is odd, we obtain  $\sum_{i=1}^d f_i(e) \leq k - 1$  for each  $e \in E(G)$ . If  $v$  is a vertex of minimum degree, then it follows that

$$\begin{aligned} d \cdot j &= \sum_{i=1}^d j \leq \sum_{i=1}^d \sum_{e \in E_G(v)} f_i(e) \\ &= \sum_{e \in E_G(v)} \sum_{i=1}^d f_i(e) \leq \sum_{e \in E_G(v)} (k - 1) = \delta(G)(k - 1), \end{aligned}$$

and this yields to the desired bound. Assume second that  $k$  is even and  $d$  is odd. If  $e \in E(G)$  is an arbitrary edge, then  $\sum_{i=1}^d f_i(e) \leq k$ . On the left-hand side of this inequality a sum of an odd number of odd summands occurs. Therefore, it is an odd number and as  $k$  is even, we obtain  $\sum_{i=1}^d f_i(e) \leq k - 1$  for each  $e \in E(G)$ . Now the desired bound follows as above, and the proof is complete.  $\square$

The special cases  $j = k = 1$ ,  $j = 1$  and  $j = k$  of Theorem 4, Corollary 5 and Theorem 7 can be found in [1], [4] and [5], respectively. According to (1),  $d_{SS}^{(j,k)}(G)$  is a positive integer. If we suppose in the case  $j = k = 1$  that  $d_{SS}(G) = d_{SS}^{(1,1)}(G)$  is an even integer, then Theorem 7 leads to the contradiction  $d_{SS}(G) \leq 0$ . Consequently, we obtain the next known result.

**Corollary 8** ([1]). *The signed star domatic number  $d_{SS}(G)$  is an odd integer.*

**Proposition 9.** *Let  $j, k$  be two integers such that  $j \geq 1$  and  $k \geq 2$ , and let  $G$  be a graph with minimum degree  $\delta(G) \geq j$ . Then  $d_{SS}^{(j,k)}(G) = 1$  if and only if each edge  $e \in E(G)$  has an endpoint  $u$  such that  $d(u) = j$  or  $d(u) = j + 1$ .*

*Proof.* Assume that each edge  $e \in E(G)$  has an endpoint  $u$  such that  $d(u) = j$  or  $d(u) = j + 1$ . It follows from Observation 1 that  $\gamma_{jSS}(G) = m$  and thus  $d_{SS}^{(j,k)}(G) = 1$ .

Conversely, assume that  $d_{SS}^{(j,k)}(G) = 1$ . If  $G$  contains an edge  $e = uv$  such that  $d(u) \geq j + 2$  and  $d(v) \geq j + 2$ , then the functions  $f_i: E(G) \rightarrow \{-1, 1\}$  such that  $f_1(x) = 1$  for each  $x \in E(G)$  and  $f_2(e) = -1$  and  $f_2(x) = 1$  for each edge  $x \in E(G) \setminus \{e\}$  are signed star  $j$ -dominating functions on  $G$  such that  $f_1(x) + f_2(x) \leq 2 \leq k$  for each edge  $x \in E(G)$ . Thus  $\{f_1, f_2\}$  is a signed star  $(j, k)$ -dominating family on  $G$ , a contradiction to  $d_{SS}^{(j,k)}(G) = 1$ .  $\square$

The next result is an immediate consequence of Observation 1 and Proposition 9.

**Corollary 10.** *Let  $j, k$  be two integers such that  $j \geq 1$  and  $k \geq 2$ , and let  $G$  be a graph with minimum degree  $\delta(G) \geq j$ . Then  $d_{SS}^{(j,k)}(G) = 1$  if and only if  $\gamma_{jSS}(G) = m$ .*

Next we present a lower bound on the signed star  $(j, k)$ -domatic number.

**Proposition 11.** *Let  $j, k$  be two integers such that  $k \geq j \geq 1$ , and let  $G$  be a graph with minimum degree  $\delta(G) \geq j$ . If  $G$  contains a vertex  $v \in V(G)$  such that all vertices of  $N[N[v]]$  have degree at least  $j + 2$ , then  $d_{SS}^{(j,k)}(G) \geq j$ .*

*Proof.* Let  $\{u_1, u_2, \dots, u_j\} \subset N(v)$ . The hypothesis that all vertices of  $N[N[v]]$  have degree at least  $j + 2$  implies that the functions  $f_i: E(G) \rightarrow \{-1, 1\}$  such that  $f_i(vu_i) = -1$  and  $f_i(x) = 1$  for each edge  $x \in E(G) \setminus \{vu_i\}$  are signed star  $j$ -dominating functions on  $G$  for  $i \in \{1, 2, \dots, j\}$ . Since  $f_1(x) + f_2(x) + \dots + f_j(x) \leq j \leq k$  for each edge  $x \in E(G)$ , we observe that  $\{f_1, f_2, \dots, f_j\}$  is a signed star  $(j, k)$ -dominating family on  $G$ , and Proposition 11 is proved.  $\square$

**Corollary 12.** *Let  $j, k$  be two integers such that  $k \geq j \geq 1$ . If  $G$  is a graph of minimum degree  $\delta(G) \geq j + 2$ , then  $d_{SS}^{(j,k)}(G) \geq j$ .*

**Corollary 13.** *Let  $j, k \geq 1$  be integers, and let  $G$  be an  $r$ -regular graph with  $r \geq j$ .*

- (1) *If  $j \leq r \leq j + 1$ , then  $d_{SS}^{(j,k)}(G) = 1$ .*
- (2) *If  $r = j + 2p + 1$  with an integer  $p \geq 1$  and  $k \geq j$ , then  $j \leq d_{SS}^{(j,k)}(G) \leq \frac{kr}{j+1}$ .*
- (3) *If  $r = j + 2p$  with an integer  $p \geq 1$  and  $k \geq j$ , then  $j \leq d_{SS}^{(j,k)}(G) \leq \frac{kr}{j}$ .*

*Proof.* (1) Assume that  $j \leq r \leq j + 1$ . According to Observation 1,  $\gamma_{jSS}(G) = m$  and thus  $d_{SS}^{(j,k)}(G) = 1$ .

(2) Assume that  $r = j + 2p + 1$  with  $p \geq 1$ . The condition  $k \geq j$  and Corollary 12 imply that  $j \leq d_{SS}^{(j,k)}(G)$ . If  $j$  is even, then  $r = j + 2p + 1$  is odd, and if  $j$  is odd, then  $r = j + 2p + 1$  is even. Therefore, Corollary 5 leads to the desired upper bound of  $d_{SS}^{(j,k)}(G)$ .

(3) Assume that  $r = j + 2p$  with  $p \geq 1$ . The condition  $k \geq j$  and Corollary 12 imply that  $j \leq d_{SS}^{(j,k)}(G)$ . In addition, Theorem 2 yields the desired upper bound of  $d_{SS}^{(j,k)}(G)$ .  $\square$

### 3. BOUNDS ON THE PRODUCT AND THE SUM OF $\gamma_{jSS}(G)$ AND $d_{SS}^{(j,k)}(G)$

Note that  $\gamma_{jSS}(G) = m$  implies immediately  $d_{SS}^{(j,k)}(G) = 1$ , and so  $\gamma_{jSS}(G) \cdot d_{SS}^{(j,k)}(G) = m$  and  $\gamma_{jSS}(G) + d_{SS}^{(j,k)}(G) = m + 1$ . In this section, we present general bounds of the product and the sum of  $\gamma_{jSS}(G)$  and  $d_{SS}^{(j,k)}(G)$ .

**Theorem 14.** *Let  $j, k \geq 1$  be integers. If  $G$  is a graph of size  $m$  and minimum degree  $\delta(G) \geq j$ , then*

$$\gamma_{jSS}(G) \cdot d_{SS}^{(j,k)}(G) \leq mk.$$

Moreover, if  $\gamma_{jSS}(G) \cdot d_{SS}^{(j,k)}(G) = mk$ , then for each  $d_{SS}^{(j,k)}$ -family  $\{f_1, f_2, \dots, f_d\}$  of  $G$ , each function  $f_i$  is a  $\gamma_{jSS}(G)$ -function and  $\sum_{i=1}^d f_i(e) = k$  for all  $e \in E(G)$ .

*Proof.* If  $\{f_1, f_2, \dots, f_d\}$  is a signed star  $(j, k)$ -dominating family on  $G$  such that  $d = d_{SS}^{(j,k)}(G)$ , then the definitions imply

$$\begin{aligned} d \cdot \gamma_{jSS}(G) &= \sum_{i=1}^d \gamma_{jSS}(G) \leq \sum_{i=1}^d \sum_{e \in E(G)} f_i(e) \\ &= \sum_{e \in E(G)} \sum_{i=1}^d f_i(e) \leq \sum_{e \in E(G)} k = mk \end{aligned}$$

as desired.

If  $\gamma_{jSS}(G) \cdot d_{SS}^{(j,k)}(G) = mk$ , then the two inequalities occurring in the proof become equalities. Hence for the  $d_{SS}^{(j,k)}$ -family  $\{f_1, f_2, \dots, f_d\}$  of  $G$  and for each  $i$ ,  $\sum_{e \in E(G)} f_i(e) = \gamma_{jSS}(G)$ , thus each function  $f_i$  is a  $\gamma_{jSS}(G)$ -function and  $\sum_{i=1}^d f_i(e) = k$  for all  $e \in E(G)$ .  $\square$

**Theorem 15.** *Let  $j, k \geq 1$  be integers. If  $G$  is a graph of size  $m$  and minimum degree  $\delta(G) \geq j$ , then*

$$d_{SS}^{(j,k)}(G) + \gamma_{jSS}(G) \leq mk + 1.$$

*Proof.* According to Theorem 14, we have

$$d_{SS}^{(j,k)}(G) + \gamma_{jSS}(G) \leq d_{SS}^{(j,k)}(G) + \frac{km}{d_{SS}^{(j,k)}(G)}.$$

Using the fact that the function  $g(x) = x + (km)/x$  is decreasing for  $1 \leq x \leq \sqrt{km}$  and increasing for  $\sqrt{km} \leq x \leq km$ , we obtain

$$d_{SS}^{(j,k)}(G) + \gamma_{jSS}(G) \leq \max \left\{ 1 + mk, mk + \frac{km}{km} \right\} = mk + 1.$$

□

Next we improve Theorem 15 considerably.

**Theorem 16.** *Let  $j, k \geq 1$  be two integers. If  $G$  is a graph of size  $m$  and minimum degree  $\delta(G) \geq j$ , then*

$$\gamma_{jSS}(G) + d_{SS}^{(j,k)}(G) \leq \begin{cases} m + 1 & \text{if } k = 1, \\ \frac{mk}{2} + 2 & \text{if } k \geq 2. \end{cases}$$

*Proof.* If  $k = 1$ , then Theorem 15 leads to the desired bound. Therefore we assume next that  $k \geq 2$ . If the order  $n = 2$ , then  $\gamma_{jSS}(G) = m = 1$  and  $d_{SS}^{(j,k)}(G) = 1$  and hence the desired bound is valid. Now we assume that  $n \geq 3$ . Let  $f$  be a SSjDF on  $G$ . Since  $\sum_{e \in E_G(v)} f(e) \geq j$  for every vertex  $v$  of  $G$ , it follows that

$$2 \sum_{e \in E(G)} f(e) = \sum_{v \in V(G)} \sum_{e \in E_G(v)} f(e) \geq \sum_{v \in V(G)} j = nj.$$

This implies  $\gamma_{jSS}(G) \geq nj/2$ . As  $n \geq 3$  and  $j \geq 1$ , we obtain  $\gamma_{jSS}(G) \geq 2$ . Theorem 14 implies that

$$\gamma_{jSS}(G) + d_{SS}^{(j,k)}(G) \leq \gamma_{jSS}(G) + \frac{mk}{\gamma_{jSS}(G)}.$$

If we define  $x = \gamma_{jSS}(G)$  and  $g(x) = x + (mk)/x$  for  $x > 0$ , then because  $2 \leq \gamma_{jSS}(G) \leq m$ , we have to determine the maximum of the function  $g$  in the interval  $I : 2 \leq x \leq m$ . Using the condition  $k \geq 2$  and the fact that  $m \geq 2$ , it is easy to see that

$$\begin{aligned} \max_{x \in I} \{g(x)\} &= \max\{g(2), g(m)\} \\ &= \max \left\{ 2 + \frac{mk}{2}, m + \frac{mk}{m} \right\} \\ &= \frac{mk}{2} + 2, \end{aligned}$$

and the proof is complete. □



**Theorem 17.** *Let  $j, k \geq 1$  be two integers. If  $G$  is a graph of size  $m$ , minimum degree  $\delta(G) \geq j$  and order  $n \geq 2p + 1$  for an integer  $p \geq 1$ , then*

$$\gamma_{jSS}(G) + d_{SS}^{(j,k)}(G) \leq \begin{cases} m + k & \text{if } 1 \leq k \leq p, \\ \frac{mk}{p+1} + p + 1 & \text{if } k \geq p + 1. \end{cases}$$

*Proof.* We proceed by induction on  $p$ . Theorem 16 shows that the statement is valid for  $p = 1$ . Now let  $p \geq 2$  and assume that the statement is true for all integers  $1 \leq i \leq p - 1$ . Then the induction hypothesis implies that  $\gamma_{jSS}(G) + d_{SS}^{(j,k)}(G) \leq m + k$  for  $1 \leq k \leq p - 1$ . Thus assume next that  $k \geq p$ . The hypothesis  $n \geq 2p + 1$  leads as in the proof of Theorem 16 to

$$\gamma_{jSS}(G) \geq \frac{nj}{2} \geq \frac{(2p+1)j}{2} \geq \frac{2p+1}{2}$$

and thus  $p + 1 \leq \gamma_{jSS}(G) \leq m$ . Therefore, it follows from Theorem 14 that

$$(2) \quad \begin{aligned} \gamma_{jSS}(G) + d_{SS}^{(j,k)}(G) &\leq \gamma_{jSS}(G) + \frac{mk}{\gamma_{jSS}(G)} \\ &\leq \max \left\{ p + 1 + \frac{mk}{p+1}, m + k \right\}. \end{aligned}$$

Note that the hypothesis  $n \geq 2p + 1$  yields to  $m \geq p + 1$ .

If  $k = p$ , then we deduce from the inequality  $m \geq p + 1$  that

$$\max \left\{ p + 1 + \frac{mk}{p+1}, m + k \right\} = \max \left\{ p + 1 + \frac{mp}{p+1}, m + p \right\} = m + p.$$

If  $k \geq p + 1$ , then

$$p + 1 + \frac{mk}{p+1} \geq m + k$$

is equivalent with  $m(k - p - 1) \geq (p + 1)(k - p - 1)$ , and this inequality is valid since  $k \geq p + 1$  and  $m \geq p + 1$ . Hence the desired result follows from (2), and the proof is complete.  $\square$

#### REFERENCES

1. Atapour M., Sheikholeslami S. M., Ghameshlou A. N. and L. Volkmann, *Signed star domatic number of a graph*, Discrete Appl. Math., **158** (2010), 213–218.
2. Haynes T. W., Hedetniemi S. T. and Slater P. J., *Fundamentals of Domination in graphs*, Marcel Dekker, Inc., New York, 1998.
3. Saei R. and Sheikholeslami S. M., *Signed star  $k$ -subdomination numbers in graphs*, Discrete Appl. Math. **156** (2008), 3066–3070.
4. Sheikholeslami S. M. and Volkmann L., *Signed star  $k$ -domatic number of a graph*, Contrib. Discrete Math. **6** (2011), 20–31.
5. ———, *Signed star  $(k, k)$ -domatic number of a graph*, submitted.
6. Wang C. P., *The signed star domination numbers of the Cartesian product*, Discrete Appl. Math. **155** (2007), 1497–1505.
7. ———, *The signed  $b$ -matchings and  $b$ -edge covers of strong product graphs*, Contrib. Discrete Math. **5** (2010), 1–10.
8. Xu B., *On edge domination numbers of graphs*, Discrete Math. **294** (2005), 311–316.

9. ———, *Two classes of edge domination in graphs*, Discrete Appl. Math. **154** (2006), 1541–1546.
10. Xu B. and Li C. H., *Signed star  $k$ -domination numbers of graphs*, (Chinese) Pure Appl. Math. (Xi'an) **25** (2009), 638–641.

S. M. Sheikholeslami, Department of Mathematics & Research Group of Processing and Communication Azarbaijan Shahid Madani University Tabriz, I. R. Iran,  
*e-mail*: `s.m.sheikholeslami@azaruniv.edu`

L. Volkmann, Lehrstuhl II für Mathematik RWTH-Aachen University 52056 Aachen, Germany,  
*e-mail*: `volkm@math2.rwth-aachen.de`