

Home Page

Title Page

Contents

OSCILLATION CRITERIA FOR HALF-LINEAR PARTIAL DIFFERENTIAL EQUATIONS VIA PICONE'S IDENTITY*

NORIO YOSHIDA[†]

Abstract. A Picone's identity is established for a class of half-linear partial differential equations, and oscillation criteria are obtained by using the Picone's identity. By reducing the oscillation problem for half-linear partial differential equations to a one-dimensional oscillation problem for half-linear ordinary differential equations, we derive various oscillation results.

Key words. Picone's inequality, oscillation criteria, half-linear, partial differential equation

AMS subject classifications. 35B05

1. Introduction. Recently there has been much interest in studying the oscillatory behavior of solutions of half-linear differential equations. There are many papers (or books) dealing with oscillations of half-linear partial differential equations, see, e.g. Bognár and Došlý [2], Došlý [3, 4], Došlý and Mařík [5], Došlý and Řehák [6] Dunninger [7], Kusano, Jaroš and Yoshida [10], Mařík [13, 14] and Yoshida [15]. Picone identity plays an important role in Sturmian comparison theory and oscillation theory of differential equations. We mention the papers [1, 2, 3, 5, 7, 10, 15] which deal with Picone identity for half-linear partial differential equations. In particular, the paper [15] treats the half-linear partial

*This research was partially supported by Grant-in-Aid for Scientific Research (C)(2) (No. 16540144), The Ministry of Education, Culture, Sports, Science and Technology, Japan.

[†]Department of Mathematics, Faculty of Science, University of Toyama, Toyama, 930-8555, Japan (nori@sci.u-toyama.ac.jp).



Page 1 of 18

Go Back

Full Screen

Close

Quit

differential equation with first order term

$$\nabla \cdot (A(x)|\nabla v|^{\alpha-1}\nabla v) + (\alpha + 1)|\nabla v|^{\alpha-1}B(x) \cdot \nabla v + C(x)|v|^{\alpha-1}v = 0. \quad (*)$$

The purpose of this paper is to establish a Picone identity for the half-linear partial differential equation

$$P_\alpha[v] \equiv \sum_{i=1}^n \frac{\partial}{\partial x_i} \left((A_i(x))^2 |\nabla_A v|^{\alpha-1} \frac{\partial v}{\partial x_i} \right) + (\alpha + 1) |\nabla_A v|^{\alpha-1} B(x) \cdot \nabla_A v + C(x)|v|^{\alpha-1}v = 0 \quad (1.1)$$

and to derive oscillation results for (1.1) using the Picone identity, where $\alpha > 0$ is a constant and

$$\nabla_A v = \left(A_1(x) \frac{\partial v}{\partial x_1}, \dots, A_n(x) \frac{\partial v}{\partial x_n} \right).$$

We note that the half-linear partial differential equation (1.1) is a generalization of (*). In fact, if

$$A_1(x) = A_2(x) = \dots = A_n(x) = A(x)^{\frac{1}{\alpha+1}} \quad (A(x) > 0),$$

we see that (1.1) reduces to

$$\nabla \cdot (A(x)|\nabla v|^{\alpha-1}\nabla v) + (\alpha + 1)A(x)^{\frac{\alpha}{\alpha+1}}|\nabla v|^{\alpha-1}B(x) \cdot \nabla v + C(x)|v|^{\alpha-1}v = 0.$$

2. Picone identity. In this section we establish a Picone identity for (1.1), and obtain a sufficient condition for every solution v of (1.1) to have a zero on \bar{G} , where G is a bounded domain in \mathbb{R}^n with piecewise smooth boundary ∂G .

It is assumed that $A_i(x) \in C(\overline{G}; (0, \infty))$ ($i = 1, 2, \dots, n$), $B(x) \in C(\overline{G}; \mathbb{R}^n)$ and $C(x) \in C(\overline{G}; \mathbb{R})$.

The domain $\mathcal{D}_{P_\alpha}(G)$ of P_α is defined to be the set of all functions v of class $C^1(\overline{G}; \mathbb{R})$ with the property that $(A_i(x))^2 |\nabla_A v|^{\alpha-1} \frac{\partial v}{\partial x_i} \in C^1(G; \mathbb{R}) \cap C(\overline{G}; \mathbb{R})$ ($i = 1, 2, \dots, n$).

THEOREM 2.1 (Picone identity). *If $v \in \mathcal{D}_{P_\alpha}(G)$, $v \neq 0$ in G , then the following Picone identity holds for any $u \in C^1(G; \mathbb{R})$:*

$$\begin{aligned}
 & - \sum_{i=1}^n \frac{\partial}{\partial x_i} \left(u \varphi(u) \frac{(A_i(x))^2 |\nabla_A v|^{\alpha-1} \frac{\partial v}{\partial x_i}}{\varphi(v)} \right) \\
 & = - |\nabla_A u - u B(x)|^{\alpha+1} + C(x) |u|^{\alpha+1} \\
 & \quad + |\nabla_A u - u B(x)|^{\alpha+1} + \alpha \left| \frac{u}{v} \nabla_A v \right|^{\alpha+1} \\
 & \quad - (\alpha + 1) (\nabla_A u - u B(x)) \cdot \Phi \left(\frac{u}{v} \nabla_A v \right) \\
 & \quad - \frac{u \varphi(u)}{\varphi(v)} P_\alpha[v],
 \end{aligned} \tag{2.1}$$

where $\varphi(s) = |s|^{\alpha-1} s$ ($s \in \mathbb{R}$) and $\Phi(\xi) = |\xi|^{\alpha-1} \xi$ ($\xi \in \mathbb{R}^n$).

Proof. A direct calculation yields

$$\begin{aligned}
 & - \sum_{i=1}^n \frac{\partial}{\partial x_i} \left(u \varphi(u) \frac{(A_i(x))^2 |\nabla_A v|^{\alpha-1} \frac{\partial v}{\partial x_i}}{\varphi(v)} \right) \\
 = & - \sum_{i=1}^n \frac{\partial u}{\partial x_i} \varphi(u) \frac{(A_i(x))^2 |\nabla_A v|^{\alpha-1} \frac{\partial v}{\partial x_i}}{\varphi(v)} \\
 & - \sum_{i=1}^n u \varphi'(u) \frac{\partial u}{\partial x_i} \frac{(A_i(x))^2 |\nabla_A v|^{\alpha-1} \frac{\partial v}{\partial x_i}}{\varphi(v)} \\
 & - \sum_{i=1}^n u \varphi(u) \left(- \frac{\varphi'(v)}{\varphi(v)^2} \frac{\partial v}{\partial x_i} \right) (A_i(x))^2 |\nabla_A v|^{\alpha-1} \frac{\partial v}{\partial x_i} \\
 & - \sum_{i=1}^n u \varphi(u) \frac{\frac{\partial}{\partial x_i} \left((A_i(x))^2 |\nabla_A v|^{\alpha-1} \frac{\partial v}{\partial x_i} \right)}{\varphi(v)}.
 \end{aligned} \tag{2.2}$$

It is easy to see that

$$\sum_{i=1}^n \frac{\partial u}{\partial x_i} \varphi(u) \frac{(A_i(x))^2 |\nabla_A v|^{\alpha-1} \frac{\partial v}{\partial x_i}}{\varphi(v)} = \varphi\left(\frac{u}{v}\right) \sum_{i=1}^n (A_i(x))^2 |\nabla_A v|^{\alpha-1} \frac{\partial v}{\partial x_i} \frac{\partial u}{\partial x_i} \tag{2.3}$$

in view of the fact that $\varphi(u)/\varphi(v) = \varphi(u/v)$. Since $u\varphi'(u) = \alpha\varphi(u)$, it can be shown that

$$\begin{aligned}
 & \sum_{i=1}^n u \varphi'(u) \frac{\partial u}{\partial x_i} \frac{(A_i(x))^2 |\nabla_A v|^{\alpha-1} \frac{\partial v}{\partial x_i}}{\varphi(v)} \\
 = & \alpha \varphi\left(\frac{u}{v}\right) \sum_{i=1}^n (A_i(x))^2 |\nabla_A v|^{\alpha-1} \frac{\partial v}{\partial x_i} \frac{\partial u}{\partial x_i}.
 \end{aligned} \tag{2.4}$$

Using the identity $\varphi'(v) = \alpha(\varphi(v)/v)$, we obtain

$$\begin{aligned} & \sum_{i=1}^n u\varphi(u) \left(-\frac{\varphi'(v)}{\varphi(v)^2} \frac{\partial v}{\partial x_i} \right) (A_i(x))^2 |\nabla_A v|^{\alpha-1} \frac{\partial v}{\partial x_i} \\ &= -\alpha \frac{u}{v} \varphi\left(\frac{u}{v}\right) \sum_{i=1}^n (A_i(x))^2 |\nabla_A v|^{\alpha-1} \left(\frac{\partial v}{\partial x_i} \right)^2. \end{aligned} \quad (2.5)$$

Combining (2.2)–(2.5), we observe that

$$\begin{aligned} & -\sum_{i=1}^n \frac{\partial}{\partial x_i} \left(u\varphi(u) \frac{(A_i(x))^2 |\nabla_A v|^{\alpha-1} \frac{\partial v}{\partial x_i}}{\varphi(v)} \right) \\ &= \alpha \frac{u}{v} \varphi\left(\frac{u}{v}\right) \sum_{i=1}^n (A_i(x))^2 |\nabla_A v|^{\alpha-1} \left(\frac{\partial v}{\partial x_i} \right)^2 \\ & \quad - (\alpha + 1) \varphi\left(\frac{u}{v}\right) \sum_{i=1}^n (A_i(x))^2 |\nabla_A v|^{\alpha-1} \frac{\partial v}{\partial x_i} \frac{\partial u}{\partial x_i} \\ & \quad - \frac{u\varphi(u)}{\varphi(v)} \sum_{i=1}^n \frac{\partial}{\partial x_i} \left((A_i(x))^2 |\nabla_A v|^{\alpha-1} \frac{\partial v}{\partial x_i} \right). \end{aligned} \quad (2.6)$$

It is easily verified that

$$\begin{aligned} & \alpha \frac{u}{v} \varphi\left(\frac{u}{v}\right) \sum_{i=1}^n (A_i(x))^2 |\nabla_A v|^{\alpha-1} \left(\frac{\partial v}{\partial x_i} \right)^2 \\ &= \alpha \left| \frac{u}{v} \right|^{\alpha+1} |\nabla_A v|^{\alpha-1} \sum_{i=1}^n (A_i(x))^2 \left(\frac{\partial v}{\partial x_i} \right)^2 \\ &= \alpha \left| \frac{u}{v} \nabla_A v \right|^{\alpha+1}. \end{aligned} \quad (2.7)$$

A simple computation shows that

$$\begin{aligned}
 & -(\alpha + 1)\varphi\left(\frac{u}{v}\right)\sum_{i=1}^n(A_i(x))^2|\nabla_A v|^{\alpha-1}\frac{\partial v}{\partial x_i}\frac{\partial u}{\partial x_i} \\
 &= -(\alpha + 1)\left|\frac{u}{v}\nabla_A v\right|^{\alpha-1}\sum_{i=1}^n\left(A_i(x)\frac{\partial u}{\partial x_i}\right)\left(\frac{u}{v}A_i(x)\frac{\partial v}{\partial x_i}\right) \\
 &= -(\alpha + 1)\left|\frac{u}{v}\nabla_A v\right|^{\alpha-1}(\nabla_A u)\cdot\left(\frac{u}{v}\nabla_A v\right).
 \end{aligned} \tag{2.8}$$

Hence, combining (2.6)–(2.8) yields the following :

$$\begin{aligned}
 & -\sum_{i=1}^n\frac{\partial}{\partial x_i}\left(u\varphi(u)\frac{(A_i(x))^2|\nabla_A v|^{\alpha-1}\frac{\partial v}{\partial x_i}}{\varphi(v)}\right) \\
 &= \alpha\left|\frac{u}{v}\nabla_A v\right|^{\alpha+1}-\left(\alpha + 1\right)\left|\frac{u}{v}\nabla_A v\right|^{\alpha-1}(\nabla_A u)\cdot\left(\frac{u}{v}\nabla_A v\right) \\
 & -\frac{u\varphi(u)}{\varphi(v)}\sum_{i=1}^n\frac{\partial}{\partial x_i}\left((A_i(x))^2|\nabla_A v|^{\alpha-1}\frac{\partial v}{\partial x_i}\right).
 \end{aligned} \tag{2.9}$$

We easily obtain

$$\begin{aligned}
 & \frac{u\varphi(u)}{\varphi(v)}\left[(\alpha + 1)|\nabla_A v|^{\alpha-1}B(x)\cdot\nabla_A v\right] \\
 &= (\alpha + 1)\left|\frac{u}{v}\nabla_A v\right|^{\alpha-1}uB(x)\cdot\left(\frac{u}{v}\nabla_A v\right).
 \end{aligned} \tag{2.10}$$

Combining (2.9) and (2.10), we find that

$$\begin{aligned}
 & - \sum_{i=1}^n \frac{\partial}{\partial x_i} \left(u \varphi(u) \frac{(A_i(x))^2 |\nabla_A v|^{\alpha-1} \frac{\partial v}{\partial x_i}}{\varphi(v)} \right) \\
 = & \alpha \left| \frac{u}{v} \nabla_A v \right|^{\alpha+1} - (\alpha+1) \left| \frac{u}{v} \nabla_A v \right|^{\alpha-1} (\nabla_A u - u B(x)) \cdot \left(\frac{u}{v} \nabla_A v \right) \\
 & - \frac{u \varphi(u)}{\varphi(v)} \left[\sum_{i=1}^n \frac{\partial}{\partial x_i} \left((A_i(x))^2 |\nabla_A v|^{\alpha-1} \frac{\partial v}{\partial x_i} \right) \right. \\
 & \left. + (\alpha+1) |\nabla_A v|^{\alpha-1} B(x) \cdot \nabla_A v \right].
 \end{aligned} \tag{2.11}$$

Since

$$\frac{u \varphi(u)}{\varphi(v)} C(x) |v|^{\alpha-1} v = C(x) |u|^{\alpha+1},$$

we conclude that (2.11) is equivalent to the desired Picone identity (2.1). \square

THEOREM 2.2. *Assume that there exists a nontrivial function $u \in C^1(\overline{G}; \mathbb{R})$ such that $u = 0$ on ∂G and*

$$M_G[u] \equiv \int_G [|\nabla_A u - u B(x)|^{\alpha+1} - C(x) |u|^{\alpha+1}] dx \leq 0.$$

Then every solution $v \in \mathcal{D}_{P_\alpha}(G)$ of (1.1) must vanish at some point of \overline{G} .

Proof. Suppose to the contrary that there exists a solution $v \in \mathcal{D}_P(G)$ of (1.1) satisfying $v \neq 0$ on \overline{G} . THEOREM 2.1 implies that the Picone-type inequality (2.1) holds for the

nontrivial function u . Integrating (2.1) over G , we obtain

$$\begin{aligned}
 0 &= -M_G[u] + \int_G \left[|\nabla_A u - u B(x)|^{\alpha+1} + \alpha \left| \frac{u}{v} \nabla_A v \right|^{\alpha+1} \right. \\
 &\quad \left. - (\alpha + 1) (\nabla_A u - u B(x)) \cdot \Phi \left(\frac{u}{v} \nabla_A v \right) \right] dx \\
 &\geq \int_G \left[|\nabla_A u - u B(x)|^{\alpha+1} + \alpha \left| \frac{u}{v} \nabla_A v \right|^{\alpha+1} \right. \\
 &\quad \left. - (\alpha + 1) (\nabla_A u - u B(x)) \cdot \Phi \left(\frac{u}{v} \nabla_A v \right) \right] dx.
 \end{aligned} \tag{2.12}$$

It is easily seen that

$$\nabla_A u - u B(x) - \frac{u}{v} \nabla_A v = v \nabla_A \left(\frac{u}{v} \right) - u B(x) = v \left[\nabla_A \left(\frac{u}{v} \right) - \frac{u}{v} B(x) \right].$$

If

$$\nabla_A \left(\frac{u}{v} \right) - \frac{u}{v} B(x) \equiv 0 \quad \text{in } G,$$

then we obtain the following

$$\nabla \left(\frac{u}{v} \right) - \frac{u}{v} B_A(x) \equiv 0 \quad \text{in } G,$$

where

$$B_A(x) = \left(\frac{B_1(x)}{A_1(x)}, \dots, \frac{B_n(x)}{A_n(x)} \right).$$

It follows from a result of Jaroš, Kusano and Yoshida [9, Lemma] that

$$\frac{u}{v} = C_0 \exp h(x) \quad \text{on } \overline{G}$$

for some constant C_0 and some continuous function $h(x)$. Since $u = 0$ on ∂G , we obtain $C_0 = 0$, and hence $u \equiv 0$. This contradicts the fact that u is nontrivial, and therefore we find

$$\nabla_A u - uB(x) \not\equiv \frac{u}{v} \nabla_A v \quad \text{in } G.$$

Hence, it follows from a result of Kusano, Jaroš and Yoshida [10, Lemma 2.1] that

$$\int_G \left[|\nabla_A u - uB(x)|^{\alpha+1} + \alpha \left| \frac{u}{v} \nabla_A v \right|^{\alpha+1} - (\alpha+1) (\nabla_A u - uB(x)) \cdot \Phi \left(\frac{u}{v} \nabla_A v \right) \right] dx > 0,$$

which, combined with (2.12), yields a contradiction. The proof is complete. \square

3. Oscillation results. We consider the half-linear partial differential equation

$$P_\alpha[v] \equiv \sum_{i=1}^n \frac{\partial}{\partial x_i} \left((A_i(x))^2 |\nabla_A v|^{\alpha-1} \frac{\partial v}{\partial x_i} \right) + (\alpha+1) |\nabla_A v|^{\alpha-1} B(x) \cdot \nabla_A v + C(x) |v|^{\alpha-1} v = 0 \quad (3.1)$$

in an unbounded domain $\Omega \subset \mathbb{R}^n$, where $\alpha > 0$ is a constant, $A_i(x) \in C(\Omega; (0, \infty))$ ($i = 1, 2, \dots, n$), $B(x) \in C(\Omega; \mathbb{R})$ and $C(x) \in C(\Omega; \mathbb{R})$.

The domain $\mathcal{D}_{P_\alpha}(\Omega)$ of P_α is defined to be the set of all functions v of class $C^1(\Omega; \mathbb{R})$ with the property that $(A_i(x))^2 |\nabla_A v|^{\alpha-1} \frac{\partial v}{\partial x_i} \in C^1(\Omega; \mathbb{R})$ ($i = 1, 2, \dots, n$).

A solution $v \in \mathcal{D}_{P_\alpha}(\Omega)$ of (3.1) is said to be *oscillatory* in Ω if it has a zero in Ω_r for any $r > 0$, where

$$\Omega_r = \Omega \cap \{x \in \mathbb{R}^n; |x| > r\}.$$

THEOREM 3.1. *Assume that for any $r > 0$ there exists a bounded and piecewise smooth domain G with $\overline{G} \subset \Omega_r$. If there is a nontrivial function $u \in C^1(\overline{G}; \mathbb{R})$ such that $u = 0$ on ∂G and $M_G[u] \leq 0$, where M_G is defined in Theorem 2.2, then every solution $v \in \mathcal{D}_{P_\alpha}(\Omega)$ of (3.1) is oscillatory in Ω .*

Proof. Let $r > 0$ be an arbitrary number. THEOREM 2.2 implies that every solution $v \in \mathcal{D}_{P_\alpha}(\Omega)$ of (3.1) has a zero on $\overline{G} \subset \Omega_r$, that is, every solution v of (3.1) is oscillatory in Ω . \square

LEMMA 3.2. *Let $0 < \alpha < 1$. Then we obtain the inequality*

$$|\nabla u - uW(x)|^{\alpha+1} \leq \frac{|\nabla u|^{\alpha+1}}{1-\alpha} + \frac{|W(x)|^{\alpha+1}}{1-\alpha} |u|^{\alpha+1} \quad (3.2)$$

for any function $u \in C^1(G; \mathbb{R})$ and any n -vector function $W(x) \in C(G; \mathbb{R})$.

Proof. The following inequality holds:

$$\begin{aligned} (\nabla u) \cdot \Phi(\nabla u) + \alpha (\nabla u - uW(x)) \cdot \Phi(\nabla u - uW(x)) \\ - (\alpha + 1) (\nabla u) \cdot \Phi(\nabla u - uW(x)) \geq 0 \end{aligned}$$

(see, e.g., Kusano, Jaroš and Yoshida [10, Lemma 2.1]). Hence, we have

$$\begin{aligned} (\nabla u) \cdot \Phi(\nabla u) + \alpha (\nabla u - uW(x)) \cdot \Phi(\nabla u - uW(x)) \\ - (\alpha + 1) (\nabla u - uW(x) + uW(x)) \cdot \Phi(\nabla u - uW(x)) \geq 0, \end{aligned}$$

and therefore

$$\begin{aligned} |\nabla u|^{\alpha+1} + \alpha |\nabla u - uW(x)|^{\alpha+1} \\ - (\alpha + 1) [|\nabla u - uW(x)|^{\alpha+1} + uW(x) \cdot \Phi(\nabla u - uW(x))] \geq 0, \end{aligned}$$

or

$$|\nabla u|^{\alpha+1} - (\alpha + 1) uW(x) \cdot \Phi(\nabla u - uW(x)) \geq |\nabla u - uW(x)|^{\alpha+1}. \quad (3.3)$$

Using Schwarz's inequality and Young's inequality, we find that

$$\begin{aligned}
 & |(\alpha + 1)uW(x) \cdot \Phi(\nabla u - uW(x))| \\
 & \leq (\alpha + 1)|uW(x)||\nabla u - uW(x)|^\alpha \\
 & \leq (\alpha + 1) \left[\frac{|uW(x)|^{\alpha+1}}{\alpha + 1} + \frac{|\nabla u - uW(x)|^{\alpha+1}}{\frac{\alpha}{\alpha}} \right] \\
 & = |uW(x)|^{\alpha+1} + \alpha|\nabla u - uW(x)|^{\alpha+1}.
 \end{aligned} \tag{3.4}$$

Combining (3.3) with (3.4) yields the following

$$\begin{aligned}
 |\nabla u - uW(x)|^{\alpha+1} & \leq |\nabla u|^{\alpha+1} + |(\alpha + 1)uW(x) \cdot \Phi(\nabla u - uW(x))| \\
 & \leq |\nabla u|^{\alpha+1} + |uW(x)|^{\alpha+1} + \alpha|\nabla u - uW(x)|^{\alpha+1},
 \end{aligned}$$

and hence

$$(1 - \alpha)|\nabla u - uW(x)|^{\alpha+1} \leq |\nabla u|^{\alpha+1} + |W(x)|^{\alpha+1}|u|^{\alpha+1},$$

which is equivalent to (3.2). The proof is complete. \square

THEOREM 3.3. *Let $0 < \alpha < 1$. Assume that for any $r > 0$ there exist a bounded and piecewise smooth domain G with $\bar{G} \subset \Omega_r$ and a nontrivial function $u \in C^1(\bar{G}; \mathbb{R})$ such that $u = 0$ on ∂G and*

$$\int_G \left[\frac{K(x)}{1 - \alpha} |\nabla u|^{\alpha+1} - \left\{ C(x) - \frac{K(x)|B_A(x)|^{\alpha+1}}{1 - \alpha} \right\} |u|^{\alpha+1} \right] dx \leq 0,$$

where $K(x) = (\max_{1 \leq i \leq n} A_i(x))^{\alpha+1}$ and

$$B_A(x) = \left(\frac{B_1(x)}{A_1(x)}, \dots, \frac{B_n(x)}{A_n(x)} \right).$$

Then every solution $v \in \mathcal{D}_{P_\alpha}(\Omega)$ of (3.1) is oscillatory in Ω .

Proof. It is easy to see that

$$|\nabla_A u - uB(x)| \leq \sqrt{\max_{1 \leq i \leq n} (A_i(x))^2} |\nabla u - uB_A(x)| \leq \left(\max_{1 \leq i \leq n} A_i(x) \right) |\nabla u - uB_A(x)|$$

and hence

$$|\nabla_A u - uB(x)|^{\alpha+1} \leq K(x) |\nabla u - uB_A(x)|^{\alpha+1}. \quad (3.5)$$

Combining (3.2) with (3.5), we obtain

$$|\nabla_A u - uB_A(x)|^{\alpha+1} \leq \frac{K(x)}{1-\alpha} |\nabla u|^{\alpha+1} + \frac{K(x)|B_A(x)|^{\alpha+1}}{1-\alpha} |u|^{\alpha+1}.$$

Therefore, we observe that

$$\begin{aligned} & \int_G [|\nabla_A u - uB(x)|^{\alpha+1} - C(x)|u|^{\alpha+1}] \, dx \\ & \leq \int_G \left[\frac{K(x)}{1-\alpha} |\nabla u|^{\alpha+1} - \left\{ C(x) - \frac{K(x)|B_A(x)|^{\alpha+1}}{1-\alpha} \right\} |u|^{\alpha+1} \right] \, dx \end{aligned}$$

and consequently, the conclusion follows from THEOREM 3.1. \square

LEMMA 3.4. *Let $E(x) \in C(G; (0, \infty))$ satisfy $E(x) > \alpha$. Then the inequality*

$$|\nabla u - uW(x)|^{\alpha+1} \leq \frac{E(x)}{E(x) - \alpha} |\nabla u|^{\alpha+1} + \frac{|E(x)W(x)|^{\alpha+1}}{E(x) - \alpha} |u|^{\alpha+1} \quad (3.6)$$

holds for any function $u \in C^1(G; \mathbb{R})$ and any n -vector function $W(x) \in C(G; \mathbb{R})$.

Proof. Proceeding as in the proof of LEMMA 3.2, we see that the inequality (3.3) holds. Applying Schwarz's inequality and Young's inequality, we have

$$\begin{aligned} & |(\alpha + 1)uW(x) \cdot \Phi(\nabla u - uW(x))| \\ &= \frac{1}{E(x)}(\alpha + 1)|uE(x)W(x)||\nabla u - uW(x)|^\alpha \\ &\leq \frac{1}{E(x)}\left(|uE(x)W(x)|^{\alpha+1} + \alpha|\nabla u - uW(x)|^{\alpha+1}\right). \end{aligned} \quad (3.7)$$

Combining (3.3) with (3.7) yields the following

$$|\nabla u - uW(x)|^{\alpha+1} \leq |\nabla u|^{\alpha+1} + \frac{|E(x)W(x)|^{\alpha+1}}{E(x)}|u|^{\alpha+1} + \frac{\alpha}{E(x)}|\nabla u - uW(x)|^{\alpha+1}$$

and therefore

$$\left(1 - \frac{\alpha}{E(x)}\right)|\nabla u - uW(x)|^{\alpha+1} \leq |\nabla u|^{\alpha+1} + \frac{|E(x)W(x)|^{\alpha+1}}{E(x)}|u|^{\alpha+1},$$

which is equivalent to (3.6). The proof is complete. \square

THEOREM 3.5. *Let $K(x) > \alpha$. Assume that for any $r > 0$ there exist a bounded and piecewise smooth domain G with $\overline{G} \subset \Omega_r$ and a nontrivial function $u \in C^1(\overline{G}; \mathbb{R})$ such that $u = 0$ on ∂G and*

$$\int_G \left[\frac{(K(x))^2}{K(x) - \alpha} |\nabla u|^{\alpha+1} - \left\{ C(x) - (K(x))^{\alpha+2} \frac{|B_A(x)|^{\alpha+1}}{K(x) - \alpha} \right\} |u|^{\alpha+1} \right] dx \leq 0.$$

Then every solution $v \in \mathcal{D}_{P_\alpha}(\Omega)$ of (3.1) is oscillatory in Ω .

Proof. We see from (3.5) and (3.6) with $E(x) = K(x)$ that

$$\begin{aligned} & \int_G [|\nabla_A u - u B(x)|^{\alpha+1} - C(x)|u|^{\alpha+1}] dx \\ & \leq \int_G \left[\frac{(K(x))^2}{K(x) - \alpha} |\nabla u|^{\alpha+1} \right. \\ & \quad \left. - \left\{ C(x) - (K(x))^{\alpha+2} \frac{|B_A(x)|^{\alpha+1}}{K(x) - \alpha} \right\} |u|^{\alpha+1} \right] dx. \end{aligned}$$

Hence, the conclusion follows from THEOREM 3.1. The proof is complete. \square

Let $\{Q(x)\}_S(r)$ denote the spherical mean of $Q(x)$ over the sphere $S_r = \{x \in \mathbb{R}^n : |x| = r\}$, that is,

$$\{Q(x)\}_S(r) = \frac{1}{\omega_n r^{n-1}} \int_{S_r} Q(x) dS = \frac{1}{\omega_n} \int_{S_1} Q(r, \theta) d\omega,$$

where ω_n is the surface area of the unit sphere S_1 and (r, θ) is the hyperspherical coordinates on \mathbb{R}^n .

THEOREM 3.6. *Let $0 < \alpha < 1$. If the half-linear ordinary differential equation*

$$\begin{aligned} & \left(r^{n-1} \left\{ \frac{K(x)}{1 - \alpha} \right\}_S (r) |y'|^{\alpha-1} y' \right)' \\ & + r^{n-1} \left\{ C(x) - \frac{K(x)|B_A(x)|^{\alpha+1}}{1 - \alpha} \right\}_S (r) |y|^{\alpha-1} y = 0 \end{aligned} \quad (3.8)$$

is oscillatory, then every solution $v \in \mathcal{D}_{P_\alpha}(\mathbb{R}^n)$ of (3.1) is oscillatory in \mathbb{R}^n .

Proof. Let $\{r_k\}$ be the zeros of a nontrivial solution $y(r)$ of (3.8) such that $r_1 < r_2 < \dots$, $\lim_{k \rightarrow \infty} r_k = \infty$. Letting

$$G_k = \{x \in \mathbb{R}^n; r_k < |x| < r_{k+1}\} \quad (k = 1, 2, \dots)$$

and $u_k(x) = y(|x|)$, we find that

$$\begin{aligned} M_{G_k}[u_k] &\leq \int_{G_k} \left[\frac{K(x)}{1-\alpha} |\nabla u_k|^{\alpha+1} - \left\{ C(x) - \frac{K(x)|B_A(x)|^{\alpha+1}}{1-\alpha} \right\} |u_k|^{\alpha+1} \right] dx \\ &= \omega_n \int_{r_k}^{r_{k+1}} \left[\left\{ \frac{K(x)}{1-\alpha} \right\}_S (r) |y'(r)|^{\alpha+1} \right. \\ &\quad \left. - \left\{ C(x) - \frac{K(x)|B_A(x)|^{\alpha+1}}{1-\alpha} \right\}_S (r) |y(r)|^{\alpha+1} \right] r^{n-1} dr \\ &= -\omega_n \int_{r_k}^{r_{k+1}} \left[\left(r^{n-1} \left\{ \frac{K(x)}{1-\alpha} \right\}_S (r) |y'(r)|^{\alpha-1} y'(r) \right)' \right. \\ &\quad \left. + r^{n-1} \left\{ C(x) - \frac{K(x)|B_A(x)|^{\alpha+1}}{1-\alpha} \right\}_S (r) |y(r)|^{\alpha-1} y(r) \right] y(r) dr \\ &= 0. \end{aligned}$$

Hence, the conclusion follows from THEOREM 3.3. \square

THEOREM 3.7. *Let $K(x) > \alpha$ in \mathbb{R}^n . If the half-linear ordinary differential equation*

$$\begin{aligned} &\left(r^{n-1} \left\{ \frac{(K(x))^2}{K(x) - \alpha} \right\}_S (r) |y'|^{\alpha-1} y' \right)' \\ &\quad + r^{n-1} \left\{ C(x) - (K(x))^{\alpha+2} \frac{|B_A(x)|^{\alpha+1}}{K(x) - \alpha} \right\}_S (r) |y|^{\alpha-1} y = 0 \end{aligned} \quad (3.9)$$



is oscillatory, then every solution $v \in \mathcal{D}_{P_\alpha}(\mathbb{R}^n)$ of (3.1) is oscillatory in \mathbb{R}^n .

Proof. The proof is quite similar to that of Theorem 3.6, and hence will be omitted. \square

Oscillation results for the half-linear ordinary differential equation

$$(p(r)|y'|^{\alpha-1}y')' + q(r)|y|^{\alpha-1}y = 0$$

have been derived by numerous authors (see, e.g., Kusano and Naito [11] and Kusano, Naito and Ogata [12]). Various oscillation results for (3.1) can be obtained by combining THEOREMS 3.6 and 3.7 with the results of [11, 12].

The following THEOREMS 3.8 and 3.9 follow by combining THEOREMS 3.3 and 3.5 with the fact that the half-linear ordinary differential equation

$$(K_0r^{n-1}|y'|^{\alpha-1}y')' + C_0r^{n-1}|y|^{\alpha-1}y = 0$$

is oscillatory for any $n \in \mathbb{N}$, $\alpha > 0$, $K_0 > 0$ and $C_0 > 0$ (see Kusano, Jaroš and Yoshida [10, Example]).

THEOREM 3.8. *Let $0 < \alpha < 1$. If there are positive constants K_0 and C_0 satisfying*

$$\frac{K(x)}{1-\alpha} \leq K_0, \quad C(x) - \frac{K(x)|B_A(x)|^{\alpha+1}}{1-\alpha} \geq C_0,$$

then every solution $v \in \mathcal{D}_{P_\alpha}(\mathbb{R}^n)$ of (3.1) is oscillatory in \mathbb{R}^n .

THEOREM 3.9. *Let $K(x) > \alpha$ in \mathbb{R}^n . If there are positive constants K_0 and C_0 satisfying*

$$\frac{(K(x))^2}{K(x) - \alpha} \leq K_0, \quad C(x) - (K(x))^{\alpha+2} \frac{|B_A(x)|^{\alpha+1}}{K(x) - \alpha} \geq C_0,$$

then every solution $v \in \mathcal{D}_{P_\alpha}(\mathbb{R}^n)$ of (3.1) is oscillatory in \mathbb{R}^n .

EXAMPLE. We consider the half-linear partial differential equation

$$\begin{aligned} & \frac{\partial}{\partial x_1} \left(|\nabla_A v| \frac{\partial v}{\partial x_1} \right) + \frac{\partial}{\partial x_2} \left(4|\nabla_A v| \frac{\partial v}{\partial x_2} \right) \\ & + 3|\nabla_A v| \left(3 \frac{\partial v}{\partial x_1} + 16 \frac{\partial v}{\partial x_2} \right) + \left(\frac{4}{3} \times 40^3 + 1 \right) |v|v = 0 \end{aligned} \quad (3.10)$$

for $x = (x_1, x_2) \in \mathbb{R}^2$, where

$$\nabla_A v = \left(\frac{\partial v}{\partial x_1}, 2 \frac{\partial v}{\partial x_2} \right).$$

Here $n = \alpha = 2$, $A_1(x) = 1$, $A_2(x) = 2$, $K(x) = 8$, $B(x) = (3, 8)$, $B_A(x) = (3, 4)$, $C(x) = (4/3) \times 40^3 + 1$. Since

$$\frac{(K(x))^2}{K(x) - \alpha} = \frac{32}{3}, \quad C(x) - (K(x))^{\alpha+2} \frac{|B_A(x)|^{\alpha+1}}{K(x) - \alpha} = 1,$$

we can take $K_0 = 32/3$ and $C_0 = 1$. It is easy to see that $K(x) > \alpha$, and hence THEOREM 3.9 implies that every solution v of (3.10) is oscillatory in \mathbb{R}^2 .

REFERENCES

- [1] W. Allegretto and Y. X. Huang, *A Picone's identity for the p -Laplacian and applications*, *Nonlinear Anal.*, **32** (1998), 819–830.
- [2] G. Bognár and O. Došlý, *The application of Picone-type identity for some nonlinear elliptic differential equations*, *Acta Math. Univ. Comenian.*, **72** (2003), 45–57.
- [3] O. Došlý, *The Picone identity for a class of partial differential equations*, *Math. Bohem.*, **127** (2002), 581–589.
- [4] O. Došlý, *Half-linear Differential Equations*, *Handbook of Differential Equations: Ordinary Differential Equations, Volume 1*, Elsevier B. V., Amsterdam, 2004.

[Home Page](#)

[Title Page](#)

[Contents](#)



Page 18 of 18

[Go Back](#)

[Full Screen](#)

[Close](#)

[Quit](#)

- [5] O. Došlý and R. Mařík, *Nonexistence of positive solutions for PDE's with p -Laplacian*, Acta Math. Hungar., **90** (2001), 89–107.
- [6] O. Došlý and P. Řehák, *Half-linear Differential Equations*, North-Holland Mathematics Studies, 202, Elsevier Science B.V., Amsterdam, 2005.
- [7] D. R. Dunninger, *A Sturm comparison theorem for some degenerate quasilinear elliptic operators*, Boll. Un. Mat. Ital. A, **9**(7) (1995), 117–121.
- [8] J. Jaroš, T. Kusano and N. Yoshida, *Picone-type inequalities for half-linear elliptic equations and their applications*, Adv. Math. Sci. Appl., **12** (2002), 709–724.
- [9] J. Jaroš, T. Kusano and N. Yoshida, *Picone-type inequalities for elliptic equations with first order terms and their applications*, J. Inequal. Appl., **2006** (2006), 1–17.
- [10] T. Kusano, J. Jaroš and N. Yoshida, *A Picone-type identity and Sturmian comparison and oscillation theorems for a class of half-linear partial differential equations of second order*, Nonlinear Anal., **40** (2000), 381–395.
- [11] T. Kusano and Y. Naito, *Oscillation and nonoscillation criteria for second order quasilinear differential equations*, Acta Math. Hungar., **76** (1997), 81–99.
- [12] T. Kusano, Y. Naito and A. Ogata, *Strong oscillation and nonoscillation of quasilinear differential equations of second order*, Differential Equations Dynam. Systems, **2** (1994), 1–10.
- [13] R. Mařík, *Hartman–Wintner type theorem for PDE with p -Laplacian*, EJTDE, **18** (2000), Proc. 6th. Coll. QTDE, 1–7.
- [14] R. Mařík, *Oscillation criteria for PDE with p -Laplacian via the Riccati technique*, J. Math. Anal. Appl., **248** (2000), 290–308.
- [15] N. Yoshida, *Oscillation of half-linear partial differential equations with first order terms*, Stud. Univ. Žilina, **17** (2003), 177–184.