

APPROXIMATE FORMULAE FOR PRICING ZERO-COUPON BONDS AND THEIR ASYMPTOTIC ANALYSIS

B. STEHLÍKOVÁ AND D. ŠEVČOVIČ

(Communicated by Lubin G. Vulkov)

Abstract. We analyze analytic approximation formulae for pricing zero-coupon bonds in the case when the short-term interest rate is driven by a one-factor mean-reverting process with a volatility nonlinearily depending on the interest rate itself. We derive the order of accuracy of the analytical approximation due to Choi and Wirjanto. We furthermore give an explicit formula for a higher order approximation and we test both approximations numerically for a class of one-factor interest rate models.

Key Words. One factor interest rate model, Cox-Ingersoll-Ross model, bond price, analytical approximation formula, experimental order of convergence.

1. Introduction

Term structure models give the dependence of time to maturity of a discount bond and its present price. One-factor models are often formulated in terms of a stochastic differential equation for the instantaneous interest rate (short rate). In the theory of nonarbitrage term structure models the bond prices (yielding the interest rates) are given by a solution to a parabolic partial differential equation. The stochastic differential equation for the short rate is specified either under a real (observed) probability measure or risk-neutral one. A risk-neutral measure is an equivalent measure such that the derivative prices (bond prices in particular) can be computed as expected values. If the short rate process is considered with a real probability measure, a function λ describing the so-called market price of risk has to be provided. The volatility part of the process is the same for both real and risk-neutral specification of the process. The changes in the drift term depend on the so called market price of risk function λ .

It is often assumed that the short rate evolves according to the following mean reverting stochastic differential equation

$$(1) \quad dr = (\alpha + \beta r)dt + \sigma r^\gamma dw$$

where $\sigma > 0$, $\gamma \geq 0$, $\alpha > 0$, β are given parameters. In particular, it includes the well known Vasicek model ($\gamma = 0$) and Cox-Ingersoll-Ross model ($\gamma = 1/2$) (c.f. Vasicek (1977) and Cox & Ross (1985)). For those particular choices of γ closed form solutions of the bond pricing PDE (2) are known. Assuming a suitable form of the market price of risk it turns out that both the real and risk neutral processes for the short rate have the form (1). More details concerning the term structure modeling can be found in Kwok (1998).

Received by the editors January 31, 2008 and, in revised form, May 22, 2008.
2000 *Mathematics Subject Classification.* 91B28, 35K05.

Using US Treasury Bills data (June 1964 - December 1989), the real probability model (1) and generalized method of moments Chan et al. (1992) estimated the parameter γ at the value 1.499. This is considered to be an important contribution, as it drew attention to a more realistic form of the short rate volatility (compared to Vasicek or CIR models). Using the same US Treasury Bills data, Nowman (1997) estimated $\gamma = 1.361$ by means of Gaussian methodology. It should be noted that these estimations of γ are beyond values $\gamma = 0$ or $\gamma = \frac{1}{2}$ for which the closed form solution of the bond prices is known in an explicit form. In Treepongkaruna & Gray (2003) a model with interest rates from eight countries using generalized method of moments and quasi maximum likelihood method has been estimated. They tested the restrictions imposed by Vasicek and CIR models using the J-statistics in the generalized method of moments and likelihood ratio statistics in the quasi maximum likelihood method. In all tested cases except of one, the restrictions $\gamma = 0$ or $\gamma = \frac{1}{2}$ were rejected. Hence, the study of the bond prices for values of γ different from 0 and 1/2 can be justified by empirical results. However, in these cases no closed form expression for bond prices is known. An approximate analytical solution was suggested in Choi & Wirjanto (2007) which could make the models with general $\gamma > 0$ to be more widely used. In this paper, we analyze the analytical approximation by Choi & Wirjanto (2007) and derive its accuracy order. Furthermore, by adding extra terms to it we derive an improved, higher order approximation of the bond prices.

The paper is organized as follows. In the second section, we derive the order of approximation of the analytical approximative solution from Choi & Wirjanto (2007). We derive a new, higher order accurate approximation. In the third section, we compare the two approximations with a known closed form solution from the CIR model ($\gamma = \frac{1}{2}$). In Appendix we provide a proof of uniqueness of a solution of a partial differential equation for bond pricing for the parameter range $\frac{1}{2} \leq \gamma < \frac{3}{2}$.

2. Accuracy of the analytic approximation formula for the bond price in the one-factor interest rate model

In Choi & Wirjanto (2007) the authors proposed an approximate analytical formula for the bond price in a one-factor interest rate model. They considered a model having a form (1) under the risk-neutral measure. It corresponds to the real measure process:

$$dr = (\alpha + \beta r + \lambda(t, r)\sigma r^\gamma) dt + \sigma r^\gamma dw$$

where $\lambda(t, r)$ is the so called market price of risk. For a general market price of risk function $\lambda(t, r)$, the price P of a zero-coupon bond can be obtained from a solution to the following partial differential equation:

$$(2) \quad -\partial_\tau P + \frac{1}{2}\sigma^2 r^{2\gamma} \partial_r^2 P + (\alpha + \beta r)\partial_r P - rP = 0, \quad r > 0, \quad \tau \in (0, T)$$

satisfying the initial condition $P(0, r) = 1$ for all $r > 0$ (see e.g. (Kwok, 1998, Chapter 7)).

Definition 1. *By a complete solution to (2) we mean a function $P = P(\tau, r)$ having continuous partial derivatives $\partial_\tau P$, $\partial_r P$, $\partial_r^2 P$ on $Q_T = [0, \infty) \times (0, T)$, satisfying equation (2) on Q_T , the initial condition for $r \in [0, \infty)$ and fulfilling the following growth conditions: $|P(\tau, r)| \leq M e^{-mr^\delta}$ and $|P_r(\tau, r)| \leq M$ for any $r > 0, t \in (0, T)$, where $M, m, \delta > 0$ are constants.*

It is worth to note that comparison of approximate and exact solutions is meaningful only if the uniqueness of the exact solution is guaranteed. The next theorem

gives us the uniqueness of a solution to (2) satisfying Definition 1. In order not to interrupt the discussion on approximate formulae for a solution to (2) a PDE based proof of the uniqueness of the exact solution is postponed to Appendix.

Theorem 1. *Assume $\frac{1}{2} < \gamma < \frac{3}{2}$ or $\gamma = \frac{1}{2}$ and $2\alpha \geq \sigma^2$. Then there exists a unique complete solution to (2).*

Now let us state the main result on approximation of a solution to (2) due to Choi & Wirjanto (2007). They proposed the following approximation P^{ap} for the exact solution P^{ex} :

Theorem 2. *(Choi & Wirjanto, 2007, Theorem 2) The approximate analytical solution P^{ap} is given by*

$$(3) \quad \begin{aligned} \ln P^{ap}(\tau, r) = & -rB + \frac{\alpha}{\beta}(\tau - B) + (r^{2\gamma} + q\tau) \frac{\sigma^2}{4\beta} \left[B^2 + \frac{2}{\beta}(\tau - B) \right] \\ & - q \frac{\sigma^2}{8\beta^2} \left[B^2(2\beta\tau - 1) - 2B \left(2\tau - \frac{3}{\beta} \right) + 2\tau^2 - \frac{6\tau}{\beta} \right] \end{aligned}$$

where $q(r) = \gamma(2\gamma - 1)\sigma^2 r^{2(2\gamma-1)} + 2\gamma r^{2\gamma-1}(\alpha + \beta r)$ and $B(\tau) = (e^{\beta\tau} - 1)/\beta$.

Derivation of the formula (3) is based on calculating the price as an expected value under a risk neutral measure. The tree property of conditional expectation was used and the integral appearing in the exact price was approximated to obtain a closed form approximation.

Authors furthermore showed that such an approximation coincides with the exact solution in the case of the Vasicek model. Moreover, they compared the above approximation with the exact solution of the CIR model which is also known in a closed form (c.f. Cox & Ross (1985)). Graphical and tabular description of the relative error in the bond prices has been also provided in Choi & Wirjanto (2007).

The main purpose of this paper is to derive the order of accuracy of the approximation formula (3) by estimating the difference $\ln P^{ap} - \ln P^{ex}$ of logarithms of approximative and exact solutions of the bond valuation equation (2). Then, we give an approximation formula of higher order and we analyze its order of convergence analytically and numerically.

2.1. Error estimates for the approximate analytical solution. In this part we derive the order of accuracy for the approximation derived by Choi & Wirjanto (2007).

Theorem 3. *Let P^{ap} be the approximative solution given by (3) and P^{ex} be the exact bond price given as a unique complete solution to (2). Then*

$$\ln P^{ap}(\tau, r) - \ln P^{ex}(\tau, r) = c_5(r)\tau^5 + o(\tau^5)$$

as $\tau \rightarrow 0^+$ where

$$(4) \quad \begin{aligned} c_5(r) = & -\frac{1}{120}\gamma r^{2(\gamma-2)}\sigma^2 [2\alpha^2(-1+2\gamma)r^2 + 4\beta^2\gamma r^4 - 8r^{3+2\gamma}\sigma^2 \\ & + 2\beta(1-5\gamma+6\gamma^2)r^{2(1+\gamma)}\sigma^2 + \sigma^4 r^{4\gamma} (2\gamma-1)^2(4\gamma-3) \\ & + 2\alpha r(\beta(-1+4\gamma)r^2 + (2\gamma-1)(3\gamma-2)r^{2\gamma}\sigma^2)]. \end{aligned}$$

The convergence is uniform w. r. to r on compact subintervals $[r_1, r_2] \subset \subset (0, \infty)$.

Remark 1. *The function $c_5(r)$ remains bounded as $r \rightarrow 0^+$ for the case of the CIR model in which $\gamma = 1/2$. More precisely, $\lim_{r \rightarrow 0} c_5(r) = -\frac{\sigma^2}{120}\alpha\beta$. If $1/2 < \gamma < 1$, then $c_5(r)$ becomes singular, $c_5(r) = O(r^{2(\gamma-1)})$ as $r \rightarrow 0^+$.*

Proof: Recall that the exact bond price $P^{ex}(\tau, r)$ for the model (1) is given by a solution of the PDE (2). Let us define the following auxiliary function: $f^{ex}(\tau, r) = \ln P^{ex}(\tau, r)$. Clearly, $\partial_\tau P^{ex} = P^{ex} \partial_\tau f^{ex}$, $\partial_r P^{ex} = P^{ex} \partial_r f^{ex}$ and $\partial_r^2 P^{ex} = P^{ex} \left[(\partial_r f^{ex})^2 + \partial_r^2 f^{ex} \right]$. Hence the PDE for the function f^{ex} reads as follows:

$$(5) \quad -\partial_\tau f^{ex} + \frac{1}{2} \sigma^2 r^{2\gamma} \left[(\partial_r f^{ex})^2 + \partial_r^2 f^{ex} \right] + (\alpha + \beta r) \partial_r f^{ex} - r = 0.$$

Substitution of $f^{ap} = \ln P^{ap}$ into equation (5) yields a nontrivial right-hand side $h(\tau, r)$ for the equation for the approximative solution f^{ap} :

$$(6) \quad -\partial_\tau f^{ap} + \frac{1}{2} \sigma^2 r^{2\gamma} \left[(\partial_r f^{ap})^2 + \partial_r^2 f^{ap} \right] + (\alpha + \beta r) \partial_r f^{ap} - r = h(\tau, r).$$

If we insert the approximate solution into (2) then, after long but straightforward calculations based on expansion of all terms into a Taylor series in τ we obtain:

$$(7) \quad h(\tau, r) = k_4(r) \tau^4 + k_5(r) \tau^5 + o(\tau^5)$$

where k_4 and k_5 are given by

$$(8) \quad \begin{aligned} k_4(r) &= \frac{1}{24} \gamma r^{2(\gamma-2)} \sigma^2 \left[2\alpha^2(-1+2\gamma)r^2 + 4\beta^2\gamma r^4 - 8r^{3+2\gamma}\sigma^2 \right. \\ &\quad + 2\beta(1-5\gamma+6\gamma^2)r^{2(1+\gamma)}\sigma^2 + \sigma^4 r^{4\gamma}(-3+16\gamma-28\gamma^2+16\gamma^3) \\ &\quad \left. + 2\alpha r (\beta(-1+4\gamma)r^2 + (2-7\gamma+6\gamma^2)r^{2\gamma}\sigma^2) \right], \\ k_5(r) &= \frac{\gamma\sigma^2}{120} r^{2(-2+\gamma)} \left[6\alpha^2\beta(-1+2\gamma)r^2 + 12\beta^3\gamma r^4 - 10(1-2\gamma)^2 r^{1+4\gamma} \sigma^4 \right. \\ &\quad + 6\beta^2\sigma^2 (1-5\gamma+6\gamma^2) r^{2(1+\gamma)} \\ &\quad + \beta r^{2\gamma}\sigma^2 \left(-10(5+2\gamma)r^3 + 3(1-2\gamma)^2(-3+4\gamma)r^{2\gamma}\sigma^2 \right) \\ &\quad + 2\alpha r \left(3\beta^2(-1+4\gamma)r^2 + 3\beta(2-7\gamma+6\gamma^2)r^{2\gamma}\sigma^2 \right. \\ &\quad \left. \left. - 5(-1+2\gamma)r^{1+2\gamma}\sigma^2 \right) \right]. \end{aligned}$$

Let us consider a function $g(\tau, r) = f^{ap} - f^{ex}$. As $(\partial_r g)^2 = (\partial_r f^{ap})^2 - (\partial_r f^{ex})^2 - 2\partial_r f^{ex} \partial_r g$ we have

$$\begin{aligned} -\partial_\tau g &+ \frac{1}{2} \sigma^2 r^{2\gamma} \left[(\partial_r g)^2 + (\partial_r^2 g) \right] + (\alpha + \beta r) \partial_r g \\ &= \left\{ -\partial_\tau f^{ap} + \frac{1}{2} \sigma^2 r^{2\gamma} \left[(\partial_r f^{ap})^2 + \partial_r^2 f^{ap} \right] + (\alpha + \beta r) \partial_r f^{ap} \right\} \\ &\quad - \left\{ -\partial_\tau f^{ex} + \frac{1}{2} \sigma^2 r^{2\gamma} \left[(\partial_r f^{ex})^2 + (\partial_r^2 f^{ex}) \right] + (\alpha + \beta r) \partial_r f^{ex} \right\} \\ &\quad - \sigma^2 r^{2\gamma} \partial_r f^{ex} \partial_r g. \end{aligned}$$

It follows from (5) and (6) that the function g satisfies the following PDE: we obtain a PDE for the function g :

$$(10) \quad \begin{aligned} -\partial_\tau g &+ \frac{1}{2} \sigma^2 r^{2\gamma} \left[(\partial_r g)^2 + \partial_r^2 g \right] + (\alpha + \beta r) \partial_r g \\ &= h(\tau, r) - \sigma^2 r^{2\gamma} (\partial_r f^{ex})(\partial_r g), \end{aligned}$$

where $h(\tau, r)$ satisfies (7). Let us expand the solution of (10) into a Taylor series with respect to τ with coefficients depending on r . We obtain $g(\tau, r) =$

$\sum_{i=0}^{\infty} c_i(r) \tau^i = \sum_{i=\omega}^{\infty} c_i(r) \tau^i$, i.e. the first nonzero term in the expansion is $c_{\omega}(r) \tau^{\omega}$. Then $\partial_{\tau} g = \omega c_{\omega}(r) \tau^{\omega-1} + o(\tau^{\omega-1})$ and $h(\tau, r) = k_4(r) \tau^4 + o(\tau^4)$ as $\tau \rightarrow 0^+$. Here the term $k_4(r)$ is given by (8). The remaining terms in (7) are of the order $o(\tau^{\omega-1})$ as $\tau \rightarrow 0^+$. Hence $-\omega c_{\omega}(\tau) = k_4(r) \tau^4$ from which we deduce, for $\omega = 5$, $c_5(r) = -\frac{1}{5} k_4(r)$. It means that $g(\tau, r) = \ln P^{ap}(\tau, r) - \ln P^{ex}(\tau, r) = -\frac{1}{5} k_4(r) \tau^5 + o(\tau^5)$ which completes the proof. \diamond

Corollary 1. *Theorem 3 enables us to compute error in yield curves which are given by $R(\tau, r) = -\frac{\ln P(\tau, r)}{\tau}$ and relative error in bond prices.*

(1) *The error in yield curves can be expressed as*

$$R^{ap}(\tau, r) - R^{ex}(\tau, r) = -c_5(r) \tau^4 + o(\tau^4) \text{ as } \tau \rightarrow 0^+;$$

(2) *The relative error¹ of P is given by*

$$\frac{P^{ap}(\tau, r) - P^{ex}(\tau, r)}{P^{ex}(\tau, r)} = -c_5(r) \tau^5 + o(\tau^5) \text{ as } \tau \rightarrow 0^+.$$

The convergence is uniform w. r. to r on compact subintervals $[r_1, r_2] \subset \subset (0, \infty)$.

Proof: The first corollary follows from the formula for calculating yield curves. To prove the second statement we note that Theorem 3 gives $\ln P^{ap} - \ln P^{ex} = c_5(r) \tau^5 + o(\tau^5)$. Hence $P^{ap}/P^{ex} = e^{c_5(r) \tau^5 + o(\tau^5)} = 1 + c_5(r) \tau^5 + o(\tau^5)$ and therefore $\frac{P^{ap} - P^{ex}}{P^{ex}} = -c_5(r) \tau^5 + o(\tau^5)$. \diamond

Remark 2. *For the CIR model with $\gamma = 1/2$ the term $k_4(r)$ defined in (8) can be simplified to $\frac{1}{24} \sigma^2 [\alpha \beta + r(\beta^2 - 4\sigma^2)]$ and hence*

$$\ln P_{CIR}^{ap}(\tau, r) - \ln P_{CIR}^{ex}(\tau, r) = -\frac{1}{120} \sigma^2 [\alpha \beta + r(\beta^2 - 4\sigma^2)] \tau^5 + o(\tau^5)$$

as $\tau \rightarrow 0^+$ uniformly w. r. to r on compact subintervals $[r_1, r_2] \subset \subset [0, \infty)$.

2.2. Improved higher order approximation formula. It follows from (3) that the term $\ln P^{ap}(\tau, r) - c_5(r) \tau^5$ is the higher order accurate approximation of $\ln P^{ex}$ when compared to the original approximation $\ln P^{ap}(\tau, r)$ from Choi & Wirjanto (2007). Furthermore, we show, that it is even possible to compute $O(\tau^6)$ term and to obtain a new approximation $\ln P^{ap2}(\tau, r)$ such that the difference $\ln P^{ap2}(\tau, r) - \ln P^{ex}(\tau, r)$ is $o(\tau^6)$ for small values of $\tau > 0$.

Let P^{ex} be the exact bond price in the model (1). Let us define an improved approximation P^{ap2} by the formula

$$(11) \quad \ln P^{ap2}(\tau, r) = \ln P^{ap}(\tau, r) - c_5(r) \tau^5 - c_6(r) \tau^6$$

where $\ln P^{ap}$ is given by (3), $c_5(\tau)$ is given by (4) in Theorem 1 and

$$c_6(r) = \frac{1}{6} \left(\frac{1}{2} \sigma^2 r^{2\gamma} c_5''(r) + (\alpha + \beta r) c_5'(r) - k_5(r) \right)$$

where c_5' and c_5'' stand for the first and second derivative of $c_5(r)$ w. r. to r and k_5 is defined in (9).

Theorem 4. *The difference between the higher order approximation $\ln P^{ap2}$ given by (11) and the exact solution $\ln P^{ex}$ satisfies $\ln P^{ap2}(\tau, r) - \ln P^{ex}(\tau, r) = o(\tau^6)$ as $\tau \rightarrow 0^+$. The convergence is uniform w. r. to r on compact subintervals $[r_1, r_2] \subset \subset (0, \infty)$.*

¹This is referred to as the relative mispricing in Choi & Wirjanto (2007)

Proof: We have to prove that $g(\tau, r) = c_5(r)\tau^5 + c_6(r)\tau^6 + o(\tau^6)$ where c_5 and c_6 are given above. We already know the form of the coefficient $c_5 = c_5(r)$. Consider the following Taylor series expansions:

$$g(\tau, r) = \sum_{i=5}^{\infty} c_i(r)\tau^i, \quad h(\tau, r) = \sum_{i=4}^{\infty} k_i(r)\tau^i, \quad f(\tau, r) = \sum_{i=1}^{\infty} l_i(r)\tau^i.$$

The absolute term l_0 is zero because $f^{ex}(0, r) = \ln P^{ex}(0, r) = \ln 1 = 0$ for all $r > 0$. Substituting power series into equation (10) and comparing coefficients of the order τ^5 enables us to derive the identity: $-6c_6(r) + \frac{1}{2}\sigma^2r^{2\gamma}c_5''(r) + (\alpha + \beta r)c_5'(r) - k_5(r) = 0$ and hence $c_6(r) = \frac{1}{6}(\frac{1}{2}\sigma^2r^{2\gamma}c_5''(r) + (\alpha + \beta r)c_5'(r) - k_5(r))$. The term $k_5(r)$ given by (9) is obtained by computing the expansion of h . \diamond

The order of relative error of bond prices and order of error of interest rates for the new higher order approximation can be derived similarly as in Corollary 1.

Remark 3. *It is not obvious how to obtain the next higher order terms of expansion because the equations contain unknown coefficients $l_i(r)$, $i \geq 1$, of logarithm of the exact solution which is not known explicitly.*

Remark 4. *In the case of the CIR model we have*

$$c_5^{CIR}(r) = -\frac{\sigma^2}{120}(\alpha\beta + r(\beta^2 - 4\sigma^2)), \quad k_5^{CIR}(r) = \frac{\beta\sigma^2}{40}(\alpha\beta + (\beta^2 - 10\sigma^2)r)$$

and so $c_6^{CIR}(r) = \frac{\sigma^2}{360}(-2\alpha\beta^2 + 17\beta\sigma^2r - 2\beta^3r + 2\alpha\sigma^2)$. Hence

$$\begin{aligned} \ln P_{CIR}^{ap2} = \ln P_{CIR}^{ap} &+ \frac{\sigma^2}{120}(\alpha\beta + r(\beta^2 - 4\sigma^2))\tau^5 \\ &- \frac{\sigma^2}{360}(-2\alpha\beta^2 + 17\beta\sigma^2r - 2\beta^3r + 2\alpha\sigma^2)\tau^6 \end{aligned}$$

The theorem yields $\ln P_{CIR}^{ap2}(\tau, r) - \ln P_{CIR}^{ex}(\tau, r) = o(\tau^6)$. By computing the expansions of both exact and this approximative solutions we finally obtain

$$\begin{aligned} \ln P_{CIR}^{ap2}(\tau, r) &= \ln P_{CIR}^{ex}(\tau, r) - \frac{\sigma^2}{5040} \left(11\alpha\beta^3 + 11\beta^4r - 34\alpha\beta\sigma^2 \right. \\ &\quad \left. - 180\beta^2r\sigma^2 + 34r\sigma^4 \right) \tau^7 + o(\tau^7) \quad \text{as } \tau \rightarrow 0^+. \end{aligned}$$

2.3. Comparison of approximations to the exact solution for the CIR model. In this section we present a comparison of the original and improved approximations in the case of the CIR model where the exact solution is known. We use the parameter values from Choi & Wirjanto (2007), i.e. $\alpha = 0.00315$, $\beta = -0.0555$ and $\sigma = 0.0894$.

In Table 1 we show L_∞ and L_2 -norms with respect to r of the difference $\ln P^{ap} - \ln P^{ex}$ and $\ln P^{ap2} - \ln P^{ex}$ where we considered $r \in [0, 0.15]$. Maximum value considered 0.15 means 15 percent interest rate, which should be sufficient for practical use. We also compute the experimental order of convergence (EOC) in these norms. Recall that the experimental order of convergence gives an approximation of the exponent α of expected power law estimate for the error $\|\ln P^{ap}(\tau, \cdot) - \ln P^{ex}(\tau, \cdot)\| = O(\tau^\alpha)$ as $\tau \rightarrow 0^+$. The EOC_i is given by a ratio

$$EOC_i = \frac{\ln(err_i/err_{i+1})}{\ln(\tau_i/\tau_{i+1})} \quad \text{where } err_i = \|\ln P^{ap}(\tau_i, \cdot) - \ln P^{ex}(\tau_i, \cdot)\|_p.$$

In Table 2 and Figure 1 we show the L_2 -error of the difference between the original and improved approximations for larger values of τ . It turned out that the

TABLE 1. The L_∞ and L_2 – errors for the original $\ln P_{CIR}^{ap}$ and improved $\ln P_{CIR}^{ap2}$ approximations


τ	$\ \ln P^{ap} - \ln P^{ex}\ _\infty$	EOC	$\ \ln P^{ap2} - \ln P^{ex}\ _\infty$	EOC
1	2.774×10^{-7}	4.930	4.682×10^{-10}	7.039
0.75	6.717×10^{-8}	4.951	6.181×10^{-11}	7.029
0.5	9.023×10^{-9}	4.972	3.576×10^{-12}	7.004
0.25	2.876×10^{-10}	–	2.786×10^{-14}	–

τ	$\ \ln P^{ap} - \ln P^{ex}\ _2$	EOC	$\ \ln P^{ap2} - \ln P^{ex}\ _2$	EOC
1	6.345×10^{-8}	4.933	9.828×10^{-11}	7.042
0.75	1.535×10^{-8}	4.953	1.296×10^{-11}	7.031
0.5	2.061×10^{-9}	4.973	7.492×10^{-13}	7.012
0.25	6.563×10^{-11}	–	5.805×10^{-15}	–

TABLE 2. The L_2 – error with respect to r for large values of τ .

τ	1	2	3	4	5
$\ \ln P^{ap} - \ln P^{ex}\ _2$	6.345×10^{-8}	1.877×10^{-6}	1.314×10^{-5}	5.093×10^{-5}	1.427×10^{-4}
$\ \ln P^{ap2} - \ln P^{ex}\ _2$	9.828×10^{-11}	1.314×10^{-8}	2.329×10^{-7}	1.799×10^{-6}	8.798×10^{-6}

τ	6	7	8	9	10
$\ \ln P^{ap} - \ln P^{ex}\ _2$	3.255×10^{-4}	6.441×10^{-4}	1.148×10^{-3}	1.890×10^{-3}	2.921×10^{-3}
$\ \ln P^{ap2} - \ln P^{ex}\ _2$	3.217×10^{-5}	9.618×10^{-5}	2.479×10^{-4}	5.705×10^{-4}	1.200×10^{-3}

FIGURE 1. The error $\|\ln P^{ap}(\tau, \cdot) - \ln P^{ex}(\tau, \cdot)\|_2$ for the original approximation (dashed line) and the new approximation (solid line). Horizontal axis is time to maturity τ .

higher order approximation P^{ap2} gives about twice better approximation of bond prices in the long time horizon up to 10 years.

2.4. Comparison of approximate and numerical solutions. In Table 3 we present a comparison of the original approximation formula with a numerical solution P^{num} . The numerical solution was obtained using a finite volume method. We used 10^5 spatial and 4.10^7 time discretization grid points in the computational domain $\tau \in [0, 1]$, $r \in [0, 0.5]$ in order to achieve the L_2 – errors less than 10^{-11} between exact solution for the CIR model and the numerical solution. The difference $O(10^{-11})$ between the numerical and approximate solutions is therefore of the same order of accuracy as the numerical scheme and hence it was not reasonable to compute EOC in this case.

TABLE 3. Norms of the difference $\ln P^{ap}(\tau, \cdot) - \ln P^{num}(\tau, \cdot)$ for several values of τ and γ .

τ	$\gamma = 0.5$		$\gamma = 0.75$	
	L_∞ norm	L_2 norm	L_∞ norm	L_2 norm
1	2.771×10^{-7}	8.967×10^{-8}	5.576×10^{-8}	1.429×10^{-8}
0.75	6.694×10^{-8}	2.165×10^{-8}	1.691×10^{-8}	3.429×10^{-9}
0.5	8.854×10^{-9}	2.867×10^{-9}	1.411×10^{-8}	4.656×10^{-10}
0.25	3.400×10^{-10}	7.236×10^{-11}	6.963×10^{-9}	9.542×10^{-11}

τ	$\gamma = 1.00$		$\gamma = 1.32$	
	L_∞ norm	L_2 norm	L_∞ norm	L_2 norm
1	5.798×10^{-9}	1.296×10^{-9}	2.664×10^{-9}	5.536×10^{-10}
0.75	1.216×10^{-9}	2.838×10^{-10}	1.406×10^{-9}	2.352×10^{-10}
0.5	9.071×10^{-10}	7.488×10^{-11}	1.113×10^{-9}	1.413×10^{-10}
0.25	6.154×10^{-10}	5.663×10^{-11}	7.860×10^{-10}	8.524×10^{-11}

3. Conclusions

We analyzed qualitative properties of the approximation formula for pricing zero coupon bonds due to Choi & Wirjanto (2007). We furthermore proposed a higher order approximation formula for pricing zero coupon bonds. We derived the order accuracy for both approximations and we test them numerically. The improved approximation is more accurate for a reasonable range of time horizons.

Acknowledgments

The authors thank the referees for their valuable comments. The support from grants DAAD-MSSR-11/2006, VEGA 1/3767/06 and UK/381/2007 is acknowledged.

Appendix A. Uniqueness of a solution to zero coupon bond PDE

In this section, we give a proof of Theorem 1. Our aim is to prove the inequality

$$(12) \quad \frac{d}{d\tau} \int_0^\infty r^\omega P^2 dr \leq K \int_0^\infty r^\omega P^2 dr$$

to be satisfied by any solution of (2) with some constants K and $\omega \geq 0$. It implies the uniqueness of a solution to the PDE (2). Indeed, if P_1 and P_2 are two solutions of (2) with the same initial condition $P(0, r) = 1$. Then $P = P_1 - P_2$ is also a solution to (2) with $P(0, r) = 0$. Let us define a function $y(\tau) = \int_0^\infty r^\omega P^2(\tau, r) dr$. Then the inequality (12) means $\frac{dy(\tau)}{d\tau} \leq Ky(\tau)$ for $\tau > 0$. It implies: $\frac{d}{d\tau} (e^{-K\tau} y(\tau)) = -Ke^{-K\tau} y(\tau) + e^{-K\tau} \frac{dy(\tau)}{d\tau} \leq 0$. Since $y(0) = 0$ and $y(\tau) \geq 0$, it follows that $y(\tau) = 0$ for all τ . Thereof $P(\tau, r) = 0$ for all $\tau \geq 0$, $r \geq 0$ and hence $P_1 \equiv P_2$ as claimed.

Now let us derive inequality (12). Multiplying the equation by $r^\omega P$, where $\omega > 0$ and $2\gamma + \omega - 1 > 0$ using the identity $\frac{1}{2} \frac{d}{d\tau} \int_0^\infty r^\omega P^2 dr = \int_0^\infty r^\omega P \partial_\tau P dr$, and integrating with respect to r from 0 to infinity we obtain²

$$(13) \quad \frac{1}{2} \frac{d}{d\tau} \int_0^\infty r^\omega P^2 = \frac{\sigma^2}{2} \int_0^\infty r^{2\gamma+\omega} \partial_r^2 PP + \int_0^\infty (\alpha + \beta r) r^\omega \partial_r PP - \int_0^\infty r^{\omega+1} P^2.$$

²In what follows, we shall omit the differential dr from the notation

We use the notation $P' = \partial_r P$, $P'' = \partial_r^2 P$. Firstly, we use integration by parts for the following integrals from the above equation:

$$\begin{aligned}\int_0^\infty r^{2\gamma+\omega} P'' P &= -(2\gamma + \omega) \int_0^\infty r^{2\gamma+\omega-1} P P' - \int_0^\infty r^{2\gamma+\omega} (P')^2 \\ &= \frac{1}{2}(2\gamma + \omega)(2\gamma + \omega - 1) \int_0^\infty r^{2\gamma+\omega-2} P^2 - \int_0^\infty r^{2\gamma+\omega} (P')^2\end{aligned}$$

where we have used the identity $\int_0^\infty r^{\omega+\xi} P' P = -\frac{\omega+\xi}{2} \int_0^\infty r^{\omega+\xi-1} P^2$ valid for any $\omega, \xi \geq 0$ and a function P satisfying the decay estimates from Definition 1. Substituting this to (13), we end up with the identity

$$\begin{aligned}\frac{1}{2} \frac{d}{d\tau} \int_0^\infty r^\omega P^2 &= \frac{\sigma^2}{4}(2\gamma + \omega)(2\gamma + \omega - 1) \int_0^\infty r^{2\gamma+\omega-2} P^2 - \frac{\sigma^2}{2} \int_0^\infty r^{2\gamma+\omega} (P')^2 \\ (14) \quad &- \frac{\alpha\omega}{2} \int_0^\infty r^{\omega-1} P^2 - \frac{(\omega+1)\beta}{2} \int_0^\infty r^\omega P^2 - \int_0^\infty r^{\omega+1} P^2.\end{aligned}$$

Case 1: $\gamma = \frac{1}{2}$ and $2\alpha \geq \sigma^2$. We recall that the condition $2\alpha \geq \sigma^2$ in the case of CIR model ($\gamma = \frac{1}{2}$) is very well understood as it almost surely guarantees the strict positivity of the stochastic processes $r = r_t$ satisfying the stochastic differential equation: $dr = (\alpha + \beta r) dt + \sigma \sqrt{r} dw$ (see e.g. Kwok (1998)).

Subcase 1a: $2\alpha > \sigma^2$. We use the equality (14) with $\gamma = 1/2$ and $\omega = \frac{2\alpha}{\sigma^2} - 1 > 0$ to obtain the desired inequality (12) with $K = (\omega+1)\beta$.

Subcase 1b: $2\alpha = \sigma^2$. Using identity (14) with $\omega = 0$ (or simply by multiplying the PDE with P and integrating over $(0, \infty)$) we obtain the inequality (12) with $K = \beta$.

Case 2: $\gamma \in (\frac{1}{2}, 1)$. We use equation (13) with $\omega = 2$ and estimate the integral $\int_0^\infty r^{2\gamma} P^2$ by using Hölder's inequality:

$$\int_0^\infty r^{2\gamma} P^2 = \int_0^\infty (r^{4\gamma-2} P^{4\gamma-2}) (r^{2-2\gamma} P^{4-4\gamma}) \leq \left(\int_0^\infty r^2 P^2 \right)^{2\gamma-1} \left(\int_0^\infty r P^2 \right)^{2-2\gamma}.$$

It follows from the Young's inequality $ab \leq \frac{1}{p\varepsilon^p} a^p + \frac{1}{q} \varepsilon^q b^q$ for $p, q \geq 1$ such that $\frac{1}{p} + \frac{1}{q} = 1$ and any $\varepsilon > 0$ we get

$$\int_0^\infty r^{2\gamma} P^2 \leq (2\gamma - 1) \left(\frac{1}{\varepsilon} \right)^{\frac{1}{2\gamma-1}} \int_0^\infty r^2 P^2 + (2 - 2\gamma) \varepsilon^{\frac{1}{2\gamma-2}} \int_0^\infty r P^2.$$

Again using (14) with $\omega = 2$ and the above estimate we obtain

$$\begin{aligned}\frac{1}{2} \frac{d}{d\tau} \int_0^\infty r^2 P^2 &\leq \frac{\sigma^2}{2}(\gamma+1)(2\gamma+1) \int_0^\infty r^{2\gamma} P^2 - \alpha \int_0^\infty r P^2 - \frac{3\beta}{2} \int_0^\infty r^2 P^2 \\ &\leq K \int_0^\infty r^2 P^2 + \left(\sigma^2(\gamma+1)(2\gamma+1)(1-\gamma) \varepsilon^{\frac{1}{2-2\gamma}} - \alpha \right) \int_0^\infty r P^2.\end{aligned}$$

where $K = \frac{\sigma^2}{2}(\gamma+1)(2\gamma+1)(2\gamma-1) \left(\frac{1}{\varepsilon} \right)^{\frac{1}{2\gamma-1}} - \frac{3\beta}{2}$. By choosing $\varepsilon > 0$ sufficiently small such that $\sigma^2(\gamma+1)(2\gamma+1)(1-\gamma) \varepsilon^{\frac{1}{2-2\gamma}} - \alpha < 0$, we finally obtain the desired inequality $\frac{1}{2} \frac{d}{d\tau} \int_0^\infty r^2 P^2 \leq K \int_0^\infty r^2 P^2$.

Case 3: $\gamma = 1$. We again use the equation (14) with $\omega = 2$. we obtain (12) with $K = 3(2\sigma^2 - \beta)$.

Case 4: $\gamma \in (1, \frac{3}{2})$. Similarly as in the case $\frac{1}{2} < \gamma < 1$ we make use of the Hölder's inequality integral estimation:

$$\int_0^\infty r^{2\gamma} P^2 = \int_0^\infty (r^{6-4\gamma} P^{6-4\gamma}) (r^{6\gamma-6} P^{4\gamma-4}) \leq \left(\int_0^\infty r^2 P^2 \right)^{3-2\gamma} \left(\int_0^\infty r^3 P^2 \right)^{2\gamma-2}$$

and, by Young's inequality, we obtain, for any $\varepsilon > 0$,

$$\int_0^\infty r^{2\gamma} P^2 \leq (3 - 2\gamma) \left(\frac{1}{\varepsilon} \right)^{\frac{1}{3-2\gamma}} \int_0^\infty r^2 P^2 + (2\gamma - 2)\varepsilon^{\frac{1}{2\gamma-2}} \int_0^\infty r^3 P^2.$$

By (14) with $\omega = 2$ we have

$$\begin{aligned} \frac{1}{2} \frac{d}{d\tau} \int_0^\infty r^2 P^2 &\leq \frac{\sigma^2}{2} (\gamma + 1)(2\gamma + 1) \int_0^\infty r^{2\gamma} P^2 - \frac{3\beta}{2} \int_0^\infty r^2 P^2 - \int_0^\infty r^3 P^2 \\ &\leq K \int_0^\infty r^2 P^2 + \left(\sigma^2 (\gamma + 1)(2\gamma + 1)(\gamma - 1)\varepsilon^{\frac{1}{2\gamma-2}} - 1 \right) \int_0^\infty r^3 P^2. \end{aligned}$$

where $K = \frac{\sigma^2}{2} (\gamma + 1)(2\gamma + 1)(3 - 2\gamma) \left(\frac{1}{\varepsilon} \right)^{\frac{1}{3-2\gamma}} - \frac{3\beta}{2}$. By choosing $\varepsilon > 0$ sufficiently small such that $\sigma^2 (\gamma + 1)(2\gamma + 1)(\gamma - 1)\varepsilon^{\frac{1}{2\gamma-2}} - 1 < 0$ we end up with the desired inequality $\frac{1}{2} \frac{d}{d\tau} \int_0^\infty r^2 P^2 \leq K \int_0^\infty r^2 P^2$.

References

Choi, Y., & Wirjanto, T. S., An analytic approximation formula for pricing zero-coupon bonds, 2007, *Finance Research Letters*, 4, 116-126.

Chan, K. L., Karolyi, G. A., Longstaff, F. A., & Sanders, A. B., An Empirical Comparison of Alternative Models of the Short-Term Interest Rate, 1992, *Journal of Finance*, 47, 1209-1227.

Cox, J., Ingersoll, J., & Ross, S., A Theory of the Term Structure of Interest Rates, 1985, *Econometrica*, 53, 385-407.

Kwok, Y. K.: *Mathematical Models of Financial Derivatives*, New York, Heidelberg, Berlin: Springer Verlag, 1998.

Nowman, K. B., Gaussian Estimation of Single-Factor Continuous Time Models of the Term Structure of Interest Rates, 1997, *Journal of Finance*, 52, 1695-1706.

Treepongkaruna, S., & Gray, S., On the Robustness of Short Term Interest Rate Models, 2003, *Accounting and Finance*, 43, 87-121.

Vašíček, O. A., An Equilibrium Characterization of the Term Structure, 1977, *Journal of Financial Economics*, 5, 177-188.

Department of Applied Mathematics and Statistics, Comenius University, Mlynská dolina, 842 48 Bratislava, Slovakia

E-mail: {stehlikova, sevcovic}@fmph.uniba.sk

URL: <http://www.iam.fmph.uniba.sk/institute/sevcovic/>